

AUTHOR'S QUERY SHEET

Author(s): R. T. Howell et al. rhpr 249124

Article title:

Article no:

Dear Author

The following queries have arisen during the editing of your manuscript and are identified on the proofs. Unless advised otherwise, please submit all corrections using the CATS online correction form.

- AQ1 In press. Any further details?
- AQ2 Please clarify fourth author's name.
- AQ3 First five authors required for et al.
- AQ4 Please give the initial for Wingard.

UNCORRECTED PROOF

Health benefits: Meta-analytically determining the impact of well-being on objective health outcomes

RYAN T. HOWELL¹, MARGARET L. KERN², &
SONJA LYUBOMIRSKY²

¹*Department of Psychology, San Francisco State University, San Francisco, CA, USA, and*
²*Department of Psychology, University of California, Riverside, CA, USA*

(Received; in final form)

Abstract

This research synthesis integrates findings from 150 experimental, ambulatory and longitudinal studies, that tested the impact of well-being on objective health outcomes. Results demonstrated that well-being positively impacts health outcomes ($r = 0.14$). Well-being was found to be positively related to short-term health outcomes ($r = 0.15$), long-term health outcomes ($r = 0.11$), and disease or symptom control ($r = 0.13$). Results from the experimental studies demonstrated that inductions of well-being lead to healthy functioning, and inductions of ill-being lead to compromised health at similar magnitudes. Thus, the effect of subjective well-being on health is not solely due to ill-being having a detrimental impact on health, but also to well-being having a salutary impact on health. Additionally, the impact of well-being on improving health was stronger for immune system response and pain tolerance, whereas well-being was not significantly related to increases in cardiovascular and physiological reactivity. These findings point to potential biological pathways, such that well-being can directly bolster immune functioning and buffer the impact of stress.

Keywords: *Objective health outcomes, physical functioning, subjective well-being, positive affect, health processes, meta-analysis*

Increasing evidence suggests that happiness not only makes people feel good, but helps them accrue numerous advantages and rewards across multiple life domains, including work (Boehm & Lyubomirsky, in press), marriage (e.g., Marks & Fleming, 1999), and coping (e.g., Scheier et al., 1989). One of the most critical domains to explore is health. Indeed, two recent literature reviews

Correspondence: Ryan T. Howell, Ph.D., Department of Psychology, 1600 Holloway Avenue, San Francisco, CA 94132, USA. E-mail: rhowell@sfsu.edu

2 R. T. Howell et al.

summarized evidence that increased well-being is associated with improved health outcomes and lower morbidity (Lyubomirsky, King, & Diener, 2005; Pressman & Cohen, 2005). For example, happy individuals report having superior health and experiencing fewer unpleasant physical symptoms (Lyubomirsky, Tkach, & DiMatteo, 2006; Mroczek & Spiro, 2005); and higher levels of trait positive affect are associated with better quality of life for cancer patients (Ostir, Markides, Black, & Goodwin, 2000). In addition, a robust negative relation has been found between positive affect and morbidity (Pressman & Cohen, 2005).

Yet, in considering directional influences, the relation between well-being and health is undoubtedly complex. Being healthy can make people happy, and being happy can bolster health. Fortunately, experimental, ambulatory, and longitudinal studies that focus on the possible impact of well-being on objective health outcomes can help disentangle causal influences. Specifically, experimental studies determine the effects of induced positive and negative transient moods and emotions on concurrent objective health outcomes. Ambulatory studies use experience sampling methodology across several days or weeks to examine how changes in daily mood relate to health outcomes. Longitudinal studies explore whether previous levels of happiness predict future levels of physical health across more extended periods. However, researchers have not, to our knowledge, explored the nature of the well-being–health link, beyond simply reporting its magnitude. To this end, the primary goal of our meta-analysis is to synthesize the literature that investigates the possible effects of well-being on objective health status, with a focus on the moderators of this link.

Defining and measuring well-being and health

Defining well-being. The independent and predictor variables considered in this meta-analysis comprise what most researchers call “subjective well-being” (SWB; the technical term for “happiness” or simply “well-being”) or, alternatively, “life satisfaction” or “positive affect”. Happiness, life satisfaction, and positive affect are considered separable yet highly correlated constructs, and typically yield a single higher-order factor (e.g., Sheldon & Lyubomirsky, 2006; Stones & Kozma, 1980). Although these constructs are fairly heterogeneous, they are strongly related, both theoretically and empirically; thus, high ratings on life satisfaction scales and positive affect scales both indicate high well-being. For example, Watson and Clark (1994) document high correlations between average daily mood reports of positive and negative affect and trait versions of these scales (r_s from 0.48 to 0.66). Furthermore, several studies have reported that the intercorrelations typically found between various measures of trait SWB are quite large (r_s from 0.44 to 0.72; Kim, 1998; Lyubomirsky & Lepper, 1999; Suhail & Chaudhry, 2004).

Accordingly, SWB is employed here as an overarching term that comprises several related phenomena, including emotional responses (i.e., the experience of frequent positive and infrequent negative moods and emotions) and global

75 judgments of life satisfaction (Diener, 2000; Diener, Suh, Lucas, & Smith, 1999).
Furthermore, because we focus in this meta-analysis on the effects of well-being
on objective health outcomes, we use the term *well-being* for positive psychological
80 constructs that are measured (e.g., positive affect, life satisfaction, optimism) or manipulated (e.g., as part of a positive emotion induction). The term well-being is then contrasted with the term *ill-being*, which we use to refer to negative psychological constructs that are measured (e.g., negative moods, stress, anger, depression) or manipulated (e.g., as part of a negative emotion induction). The research cited used a variety of measures of the different components of well-being.

85 *Measuring well-being.* Given this article's focus on well-being's impact on health, we have included studies that use measures of both trait levels of well-being (in the longitudinal research) and transient (state level) emotions and moods (in the ambulatory and experimental research). Measures of these constructs employ self-report methods, which appropriately allow the final judge of happiness and satisfaction to be "whoever lives inside a person's skin" (Myers & Diener, 1995, p. 11; see also Diener, 1994). However, the fact that self-reports are subjective does not mean that they are unrelated to relatively more "objective" variables (for a review, see Diener, 1994). For example, research reveals significant convergence of self-reported well-being with informant reports (e.g., Lyubomirsky & Lepper, 1999; Sandvik, Diener, & Seidlitz, 1993), recall of positive and negative events (e.g., Seidlitz, Wyer, & Diener, 1997), unobtrusive observations of non-verbal (i.e., smiling) expressions (e.g., Harker & Keltner, 2001), and physiological responses (e.g., Lerner, Gonzalez, Dahl, Hariri, & Taylor, 2005).

95 Typically, longitudinal and ambulatory studies assess well-being via self-report. The longitudinal studies described here included several different measures of global well-being, including, but not limited to, the Satisfaction With Life Scale (SWLS; Diener, Emmons, Larsen, & Griffin, 1985), the Memorial University of Newfoundland Scale of Happiness (Kozma & Stones, 1980), and various single-item scales (e.g., "How satisfied are you with your life?"). We also included longitudinal studies that employed more indirect indicators of well-being, such as measures of optimism (e.g., the Life Orientation Test [LOT]; Scheier & Carver, 1985). Optimism has been found to be related to positive affectivity, and, thus, serves as a defensible proxy for well-being (Lucas, Diener, & Suh, 1996). For example, Lyubomirsky et al. (2005) found a very high correlation ($r=0.60$) between optimism and happiness. However, some specific non-hedonic quality-of-life (QOL) measures were not included, as these measures focus primarily on physical symptoms, health problems, and medical issues (e.g., QLQ-30, Aaronson et al., 1993; FACT, Cella et al., 1993; see Gotay, 2006, for a review of studies linking these types of QOL to survival) and not on emotional responses or global judgments of life satisfaction.

100
105
110
115 The ambulatory (and some experimental) studies described here typically included self-reported measures of emotions and moods, such as the Affect Balance Scale (Bradburn, 1969), variants of the Positive and Negative Affect

4 R. T. Howell et al.

Schedule (PANAS; Watson, Clark, & Tellegen, 1988), and the Profile of Mood States (Curran, Andrykowski, & Studts, 1995). Such measures are appropriate to use in ambulatory studies, whose purpose is to track small changes in affect over time; and these are the only measures available to researchers interested in measuring affect in experimental studies, as no investigations to date, to our knowledge, have tested the effects of induced long-term happiness on health. Although transient mood is not equivalent to long-term happiness, it has notably been shown to be the very *hallmark*, or basic constituent, of happiness. Indeed, happiness has been defined as the experience of frequent positive emotions (Diener, Sandvik, & Pavot, 1991; Lyubomirsky et al., 2005). Hence, we expected the physical health outcomes of short-term positive moods to be parallel to those for the concomitants of global, long-term well-being (see Lyubomirsky et al., 2005, for a similar approach).

In the experimental research reported here, a variety of manipulations were used to induce transient emotions, including films, imagery, music, and the Velten induction task (Velten, 1968), among others (see Coan & Allen, 2007, for an overview). However, many researchers induce global positive affect or positive emotions, and do not discriminate among specific emotions (e.g., happiness, elation, or arousal) or moods. Notably, positive moods are not the opposite of negative moods. These two types of affect show moderate inverse relations across individuals, sometimes correlate with different variables, and appear to be rooted in distinct biological systems (Bradburn & Caplovitz, 1965; Cacioppo, Gardner, & Berntson, 1999; Diener & Emmons, 1984; Diener, Smith, & Fujita, 1995).

Defining health. Health is a multi-dimensional construct that evades simple classification (Cacioppo & Berntson, 2007; Gochman, 1997). It can be conceptualized as two distinct categories – as a state or as a process (Carver, 2007; Kaplan, 1994, 2003) – but is usually defined as a state, extending from the traditional biomedical model. Accordingly, health is characterized by a lack of illness or disease (e.g., lack of fever, vomiting, chronic conditions, disability), maintaining normal function (ability to function well with minimal medical care), and positive self-assessments of health at the time of measurement (Breslow, 1972; Idler & Kasl, 1991). Health can be operationalized in a variety of ways, ranging from subjective single-item judgments of overall health to specific physiological measures, such as concentrations of hormones and substances in the bloodstream.

In contrast, several theorists have suggested that health be defined as a lifelong process (Aldwin, Spiro, Levenson, & Cupertino, 2001; Baltes, Staudinger, & Lindenberger, 1999; Clipp, Pavalko, & Elder, 1992; Schultz & Heckhausen, 1996); that is, health involves regulation over time, such that the autonomic, neuroendocrine, and immune systems work together to maintain balance within the body (Cacioppo & Berntson, 2007). If this balance is threatened for an extended period, these systems can break down and lead to physical decline (McEwen, 1998; McEwen & Stellar, 1993). In this meta-analysis, we define health according to this second, more holistic framework.

Measuring health. If health is considered as a process, a measurement at any single assessment reflects the individual's state within this broader process. For healthy individuals, maintaining a state of normal functioning and preventing disease are important goals, whereas for individuals with chronic illnesses, maintaining well-being and controlling symptoms are important goals (Westmas, Gil-Rivas, & Cohen Silver, 2007). Accordingly, how health is operationalized and measured depends on the person's position on the continuum between optimal functioning and clinical illness. Further, measures of health depend on whether researchers are interested in short-term markers of system activity (e.g., heart rate, blood pressure, cortisol levels) or long-term markers of overall health (e.g., cardiovascular fitness, survival). One goal of this meta-analysis is to integrate multiple levels of health; therefore, we included studies assessing both markers of physiological functioning and markers of overall functioning. Specifically, at the molecular level, health is marked by normal responses to stress and rapid recovery to baseline levels (Kemeny, 2007). At the molar level, for healthy individuals, well-being should maintain or increase normal functioning, and decrease risk of illness (such as colds and infections), and early mortality. For individuals with a chronic condition, well-being should decrease symptoms of illness (e.g., allergic reactions, asthmatic symptoms) and increase survival (longer life, despite the presence of one or more morbidities).

Finally, although health can be construed as a complex construct with multiple physical, cognitive, and affective dimensions (Fisher, 1995; Ryff & Singer, 1998; Ware, 1987), in the present analysis, we opted to operationalize health in terms of physiological measures and relatively objective physical outcomes rather than using subjective self-reports. This practice has several advantages. Shared method variance between self-reported measures of health and well-being may be responsible for a strong association between these two constructs (Lyubomirsky et al., 2005; Pressman & Cohen, 2005). Additionally, objective measures are valuable from a public health perspective in which optimal health is achieved by extending life expectancy while compressing morbidity to the final years of life (Fries, 1990; Kaplan, 2003).

Markers of normal functioning. In the literature on health and stress, several markers of hormonal responses and immune functioning are used to assess the effects of stress at the molecular level (Rabin, 1999; Segerstrom & Miller, 2004). For example, in the immune system, markers of normal immune responses include increases in lymphocytes (e.g., *t* cell counts on markers such as D4, CD8+ and CD16+), leukocytes (such as natural killer cells, macrophages, and interleukin cells), and immunoglobulin (such as sIgA) after being exposed to an invading substance (Dayyani et al., 2004; Dreher, 1995; Linnemeyer, 1993; Perera, Sabin, Nelson, & Lowe, 1998; Rabin, 1999; Saleh et al., 1995). In the autonomic nervous system (ANS), stress activates the system, evidenced by the release of specific hormones (e.g., cortisol, adrenaline) and increases in heart rate, blood pressure, finger temperature, and skin conductance (Cacioppo & Tassinari, 1990; Pickering, 1999). This response should then taper back to baseline

6 *R. T. Howell et al.*

levels. Non-normal responses may represent some sort of dysregulation within the system (Kemeny, 2007).

At the molar level, health depends on a person's status. For healthy individuals, measures of health reflect normal or optimal functioning. For example, cardiovascular strength offers a marker of the organism's level of fitness, and is commonly assessed by increased power output and flow rates (Koehler, 1996). General health indicates overall functioning, and can be measured by indicators such as a healthy cholesterol ratio and weight (Kivimäki et al., 2005; Pollard & Schwartz, 2003). Finally, longevity is a reliable and objective health outcome that is arguably the endpoint in a long causal chain of inter-related events. Longevity is determined by length of life in years, and is generally verified from vital records or familial report (e.g., Brown, Butow, Culjak, Coates, & Dunn, 2000; Friedman et al., 1993; Wingard, Berkman, & Brand, 1994).

For individuals with one or more chronic conditions, healthy functioning is marked by symptom control. For example, allergic reactions and asthma attacks indicate functional decline and dysregulation. Health is evident when a normal level of functioning can be maintained. Allergic reactions are typically measured by skin tests, in which allergens are introduced percutaneously and flare or wheel sizes are measured (Zachariae, Jorgensen, Egekvist, & Bjerring, 2001). Decreased respiratory functioning may signal respiratory failure (Quanjer et al., 1993; Quanjer, Lebowitz, Gregg, Miller, & Pedersen, 1997; Rosenow, 2005) and poor symptom control. Respiratory system functioning is commonly assessed by expiratory volume (a measure of how much air a person exhales during a forced breath), peak expiratory volume (the maximal amount of flow expelled in a forced breath), and oxygen saturation (the percentage of oxygen the red cells carry). Finally, terminal illnesses (e.g., cancer, HIV) progress through a series of stages and are marked by whether the individual declines (a lack of symptom control), maintains a stable level of functioning, or evidences some degree of recovery. Survival indicates how long a person stays alive despite having one or more chronic illness (Pressman & Cohen, 2005).

Although health investigators typically study a single illness or physiological marker of normal functioning, it is important to consider bodily systems as a whole, despite the complexity of the relations involved. For example, if well-being is indeed beneficial, then it should benefit health across systems and levels, regardless of the specific mechanisms and pathways involved (Rabin, Kusnecov, Shurin, Zhou, & Rasnick, 1994). Thus, the present meta-analysis empirically examines the potential benefits of well-being across multiple markers of normal functioning, including (a) specific, short-term outcomes, (b) general, long-term markers of physical well-being and functioning, and (c) symptom control during stages of chronic conditions. Although limited empirical support exists in human studies on the complete process connecting short-term and long-term health outcomes (Keller, Shiflett, Schleifer, & Bartlett, 1994), physiological responses may indeed extend to clinical disease outcomes over time (Keller et al., 1994; Kemeny, 2007), and by combining the two in a single analysis, we can examine

250 the macro–micro level linkages as a whole (Mroczek, Almeida, Spiro, & Pafford, 2006; Nesselrode, 1988, 1991).

Understanding the potential impact of well-being on health

Linking emotion and health. Much of the theoretical and empirical work linking psychological and physical well-being comes from studies on stress and health. Comprehensive models relating stress to health outcomes have been elaborated (e.g., Carver, 2007; Keller et al., 1994; Rabin, 1999). For example, cortisol is often used as a marker of stress. An increase in cortisol is an adaptive response to a stressor, but when prolonged over time, it can negatively impact immune system functioning (Cohen & Williamson, 1991; Dickerson & Kemeny, 2004; Herbert & Cohen, 1993; Segerstrom & Miller, 2004). Further, multiple studies and reviews indicate that stress can negatively affect the cardiovascular system (e.g., Krantz & McCeney, 2002; Kubzansky & Kawachi, 2000), neuroendocrine activity, and negative disease outcomes (Carver, 2007).

265 In a very basic model, when a physical or emotional stressor is encountered, distress occurs (Keller et al., 1994). This may, in turn, activate the central nervous system, triggering a fight-or-flight response, characterized by physiological changes, such as increased blood sugar levels, heart rate, and blood pressure, and the release of stress hormones (such as cortisol and epinephrine; Cannon, 1932; Selye, 1956). This response may directly and indirectly influence immune functioning (Glaser & Kiecolt-Glaser, 1994; Keller et al., 1994). Immune response dysregulation, in turn, may continue to activate the central nervous system, leading to chronic strain and increased susceptibility to illness (Bowen, 2001; Dickerson & Kemeny, 2004). Thus, the cardiovascular, neuroendocrine, and immune systems work together and influence one another (Cacioppo & Berntson, 2007). Some evidence for this complete model comes from animal studies with mice and non-human primates (Laudenslager & Fleshner, 1994; Maynahan et al., 1994), which suggest that such a process of dysregulation over time can potentially lead to clinical illness. Unfortunately, empirical studies that test direct and indirect pathways are mostly lacking in human research (Keller et al., 1994), although the few studies that exist offer some support (see Cohen, 1994; Keller et al., 1994, for reviews). We note that this is a simplistic description, and the system is undoubtedly much more complex (Friedman, 2007).

280 Although elements within the cardiovascular, endocrine, and immunological systems play various roles within the stress–disease process, if each system is considered as a whole, then multiple markers can be combined in an informative manner. For example, participants are often subjected to a stressor, and heart rate, blood pressure, saliva cortisol, and plasma concentrations of epinephrine are measured to determine the extent of their reactivity (e.g., Bachen et al., 1992; Kiecolt-Glaser, Malarkey, Cacioppo, & Glaser, 1994; Manuck, Cohen, Rabin, Muldoon, & Bachen, 1991). Similarly, multiple markers of immune response are often measured in response to stress. Segerstrom and Miller (2004) offer an excellent overview of the immune system, with evidence on how stress links to

285

290

8 R. T. Howell et al.

natural and specific immune responses. Although cost and participant constraints limit what physiological aspects a researcher considers (Keller et al., 1994), multiple markers may be telling a parallel story, which can be informative on how the body functions as a whole. Hence, in the present meta-analysis, we considered health in this more holistic fashion, combining multiple markers of function and dysregulation within each main system in the body and in overall functioning.

How well-being may influence health. If stress and negative emotions potentially foster detrimental health outcomes, can positive emotions and moods foster improved health? That is, whereas stress activates the sympathetic nervous system, an opposite reaction may decrease sympathetic system activity (Rabin et al., 1994), and promote optimal functioning. Empirical support for this notion is evident in personality research, which has demonstrated that negative traits, such as neuroticism and hostility, relate to increased mortality risk and poor health outcomes (e.g., Booth-Kewley & Friedman, 1987; Smith, 2006; Smith, Glazer, Ruiz, & Gallo, 2004; Smith & Williams, 1992; Suls & Bunde, 2005; Watson & Pennebaker, 1989), whereas positive traits, such as optimism, extraversion, agreeableness, and conscientiousness, relate to decreased mortality risk and better health (Friedman et al., 1993; Hampson, Goldberg, Vogt, & Dubanoski, 2006). Because of the strong correlation between personality traits and SWB, similar mechanisms may characterize the relations between well-being and health (Pressman & Cohen, 2005; Ryff & Singer, 1998).

Specifically, Pressman and Cohen (2005) detailed two models linking positive affect and disease. In the *direct effects model*, positive affect may directly affect health practices, decrease autonomic nervous system activity, regulate the release of stress hormones, influence the opioid system and immune responses, and affect social networks; these, in turn, impact health and disease outcomes. In the *stress-buffering model*, positive affect may ameliorate the effects of stressful events by increasing resiliency and enhancing coping responses. Accordingly, well-being may affect health by enhancing short-term responses (e.g., increasing immune response and pain tolerance) and long-term functioning (e.g., better cardiovascular fitness and longer life) or by buffering the effects of short-term stressors (marked by high-level stress responses and heart reactivity), and long-term illness (e.g., slowing disease progression and increasing survival). Most likely, a combination of these two mechanisms operate, depending on the individual and the situation (Friedman, 2007). In turn, health status influences well-being and quality of life.

Possible moderators of the health–SWB relation

Based on the theoretical literature and empirical studies outlined above, a primary goal of this meta-analysis was not only to establish *whether* well-being influences health outcomes, but also *under what conditions* well-being may exert its salutary effects. Accordingly, in addition to the overall relation of well-being to health, we examined several potential moderators of these relations.

335 *Categories of health outcomes.* First, we combined short- and long-term outcomes
using both general and specific markers of health. Specifically, at a broad level, we
identified three types of health outcomes according to how health can be
340 conceptualized, based on both length of follow-up and initial health status: short-
term outcomes, long-term outcomes, and disease and symptom control for
chronically ill samples. Across these three categories, we examined the effects of
12 groups of health outcomes. Specifically, short-term outcomes included (a)
immune system response, (b) cardiovascular reactivity, (c) endocrine system
functioning and response, (d) physiological response, and (e) pain tolerance.
345 Long-term outcomes included (a) general health outcomes, (b) cardiovascular
functioning, (c) respiratory functioning, and (d) longevity. Finally, disease and
symptom control included (a) measures of respiratory control (in conditions such
as allergies and asthma), (b) disease progression, and (c) survival despite having
one or more terminal conditions. In turn, each of these 12 health outcomes was
comprised of various specific markers.

350 *Health outcome as a moderator.* Our moderator predictions were based on
theories from the stress and health literature (e.g., Rabin, 1999; Segerstrom &
Miller, 2004) and the pathway models enumerated by Pressman and Cohen
(2005), as well as other relevant work. We expected well-being to relate positively
355 to *and increase* health-related functioning (i.e., longevity, survival, and pain
tolerance), improve autonomic nervous system response (i.e., cardiovascular and
respiratory functioning), and improve immune system functioning. In contrast,
we expected well-being to relate negatively to *and decrease* cardiovascular
reactivity (e.g., heart rate, blood pressure), endocrine response (e.g., measures
of cortisol), physiological response (e.g., finger temperature), symptom response
360 in chronic conditions, and disease progression. These negative relations were
predicted because well-being is expected to buffer the system from negative
outcomes. Furthermore, we predicted that well-being would affect short-term
outcomes more than long-term outcomes. When stress occurs, the autonomic
nervous system is immediately activated; if well-being interrupts this response
365 (either through buffering stress or engendering a more rapid recovery), then its
effects will be evident fairly quickly (Rabin et al., 1994). For long-term outcomes,
a vast array of variables can moderate and intercede in this relation (Hall,
Anderson, & O'Grady, 1994), ranging from psychosocial factors, such as social
support, health habits, and natural physiological changes that occur with age, to
370 measurement unreliability; hence, any effects on long-term outcomes will be
weaker, although still significant in a practical sense (Rosenthal, 1991; Smith,
2006).

375 *Baseline health as a moderator.* For healthy individuals, bodily systems naturally
fluctuate with transient stress; thus, seemingly abnormal levels of one marker may
actually be a normal response to fluctuation in another system (Rabin et al.,
1994). In contrast, for unhealthy individuals, the system as a whole is
dysregulated, and abnormal values indicate further stress on the system, adding

to the overall allostatic load (McEwen, 1998). Due to its differential role in defining and understanding health outcomes, baseline health is important to consider. We expected well-being to have a greater effect for unhealthy samples than for healthy samples.

Operationalizations of well-being. Although study methodology typically dictates the operational definition of well-being in any particular design (i.e., transient emotions are typically measured in ambulatory studies and manipulated in experimental research, whereas trait levels of well-being are typically measured in longitudinal data), we expected the relation between well-being and objective health to vary as a function of state *vs.* trait operationalizations. Specifically, we hypothesized that short-term health outcomes would be more strongly associated with state manipulations of well-being, and that long-term outcomes would be more strongly associated with trait measures of well-being.

When stress occurs and the ANS is activated (releasing cortisol, increasing heart rate and blood pressure, etc.), transient positive emotions can have relatively immediate effects – for example, potentially moderating the stress response or enabling a quicker return to baseline, indirectly protecting other systems (such as the immune system) from the stressor. Thus, much like a stressor provokes a short-term response from the ANS, positive emotions may have a short-term counteracting influence on the stressor. In contrast, long-term health outcomes represent a process of accumulated regulation or dysregulation over time. Thus, with respect to long-term outcomes, well-being that is stable over time (i.e., trait well-being) can aid individuals to maintain stability, both internally and externally, thus avoiding system dysregulation and decreasing susceptibility to illness.

Age as a moderator. In both humans and animals, the immune system changes with advancing age (Bildler, 1975; Makinodan et al., 1991; Weksler & Hausman, 1982). Specifically, a general decline in immune response may occur with age, which increases susceptibility to infections and disease (see Solomon & Benton, 1994, for a review). Notably, some factors moderate this decline. Studies with elderly individuals demonstrate that successful agers have stronger indices of immune function than normal and declining elderly individuals (Solomon et al., 1988; Thomas, Goodwin, & Goodwin, 1985), suggesting that factors other than age itself are important. Well-being may be one such factor (Kiecolt-Glaser et al., 1994). Thus, sample age is important to consider as a factor in the well-being and health relation. When there is a greater possibility of health decline, changes in health outcomes will be more evident (Rowe & Kahn, 1987; Solomon & Benton, 1994), therefore, we expected well-being to have stronger effects on health outcomes for older samples.

Gender as a moderator. As males and females differ physiologically, gender differences may impact the role that well-being plays. For example, females typically live longer than males; yet males who reach older age are often both

420 physically healthier than females, and suffer from fewer psychological problems,
such as anxiety and depression (Guralnik & Kaplan, 1989; Roos & Havens, 1991;
Strawbridge, Cohen, Shema, & Kaplan, 1996). Thus, if those males who
experience longevity also report less negative affect and more happiness, then it
is possible that well-being plays a role in the observed gender differences in health
425 outcomes for elderly samples. We predicted that well-being will be more
important for males, acting as a buffer against decline.

Previous research syntheses examining the health–SWB relation

To our knowledge, only two literature reviews to date have scrutinized the link
between well-being and health; both were published in 2005. The first was a
430 meta-analytic review of the relation of happiness and positive affect to a variety of
indicators of “success”, including health (Lyubomirsky et al., 2005); and the
second was a qualitative review that focused on the link between positive affect
(PA) and health (Pressman & Cohen, 2005).

The Lyubomirsky et al. review used three classes of evidence – cross-sectional,
435 longitudinal, and experimental – to examine the extent to which various
indicators of well-being were associated with successful outcomes (e.g., income
and marriage), and with behaviors and attributes paralleling success (e.g.,
prosocial behavior, sociability, and creativity). However, computation of effect
sizes between SWB and health outcomes constituted only a portion of the
440 analyses conducted for this review. Furthermore, Lyubomirsky and her colleagues’
analyses diverged from those of the current study in two critical ways. First,
they did not examine any moderators of the well-being–health relation (nor of
any of the well-being–success links they described). Nor did they distinguish
between the different types of health outcomes (e.g., immune functioning *vs.*
445 cardiovascular reactivity *vs.* survival) or whether those outcomes were short-term
or long-term. Their goal was simply to test whether a positive association was
present between well-being and success. Second, unlike the present study, their
analyses of health outcomes did not separate objective indicators and subjective
reports of health.

450 By contrast, the Pressman and Cohen (2005) review was a qualitative synthesis
of the PA-health literature. The authors concluded that positive affect was related
to many objective health outcomes, including lower morbidity, decreased
symptoms, and diminished reported pain, among others. Whereas they provide
great detail about the association between PA and health, our review differs from
455 their review in three important ways. First, we expanded our analyses to include
all positive psychological constructs in an attempt to determine how well-being in
general (and not only positive affect) influences objective health outcomes. We
believe this to be important, as many studies in the area of health psychology
assess the cognitive component of well-being and would have otherwise been
460 excluded. Second, we aimed to examine the differential effects of positive and
negative psychological constructs on health outcomes. To this end, we included
studies that simultaneously measured or manipulated both positive and negative

psychological constructs. Although the present meta-analysis does not provide a complete review of the effects of ill-being on health, it does afford an opportunity to compare the effect sizes for the two constructs of well-being and ill-being. Such a comparison may illuminate unique relations and pathways between health and these two constructs, as well as the mechanisms and moderators underlying them. Third, in contrast to Pressman and Cohen's qualitative review, our meta-analytic review is able to estimate the size of the effect between well-being and objective health, as well as to test quantitatively for moderators of the well-being–health link.

Objectives of the present meta-analysis

Given that numerous studies and reviews have considered the effect of negative emotions (or more generally *ill-being*) on compromised health functioning and increased illness (e.g., Friedman & Booth-Kewley, 1987; Herbert & Cohen, 1993; Segerstrom & Miller, 2004), the current meta-analysis focused on the effect of positive psychological constructs (or more generally *well-being*) on objective health outcomes. Furthermore, although several authors have recently suggested that health psychologists need to move beyond the medical model and consider health more broadly (e.g., Grzywacz & Keyes, 2004; Kaplan, 2003; Ryff & Singer, 1998; Smith & Spiro, 2002), we argue that any review of the literature must separately consider specific components of health. Thus, for the purpose of this research synthesis, we concentrated on measures of objective health outcomes using traditional biomedical markers. To these ends, our search strategy included seeking out literature examining positive emotions and positive traits as predictors of objective measures of physical and physiological health outcomes. Although the focus of the meta-analysis was on positive psychological constructs, if an included study reported the relation between well-being and health *as well as* the relation between ill-being and health, the size of the effect between ill-being and health was computed separately. This procedure allowed us to compare the average ill-being–health effect size with the average well-being–health effect size for those studies that examined both linkages. Such studies were expected to have similar effect sizes to those that examined only one of the links.

As our meta-analysis was specifically concerned with assessing the potential impact of well-being on objective health, only studies that used experimental, ambulatory, and longitudinal methods were included. The copious cross-sectional literature has been reviewed in other sources (Lyubomirsky et al., 2005; Pressman & Cohen, 2005), yet some of the studies included in those reviews suffer from multiple limitations. First, correlational and cross-sectional studies provide little information regarding directionality, as these methodologies cannot test the possible impact of well-being on health. Second, due to their common reliance on self-reports of both well-being and health, these studies generally contain too much shared method variance to determine whether well-being has any tangible impact on important health outcomes. In contrast,

particular sections of our Results pay special attention to studies that use experimental methodology, as true experiments allow us to estimate the causal effects of well-being on objective physical and physiological outcomes.

In sum, the primary aims of this meta-analysis were to (a) determine the average effect size between well-being and objective health, (b) compare this effect size to the average effect size for ill-being and health, (c) establish which particular health outcomes are most strongly associated with well-being, and (d) explore possible sample-specific moderators of the well-being–health relation.

Method

Literature search procedures

The present meta-analysis used several search techniques to retrieve all applicable studies for inclusion. Our primary search procedure extended work from the two recent reviews of health outcomes associated with positive affect and SWB (Lyubomirsky et al., 2005; Pressman & Cohen, 2005). First, each empirical article considered in these two reviews that addressed well-being and health was located; the total number of unique empirical articles examined by Lyubomirsky et al. or Pressman and Cohen was 240. Second, all literature reviews and theoretical articles cited in the two 2005 reviews was located, resulting in an additional 30 papers (for a total of 270) to be used for additional search techniques (i.e., forward and backward searching).

Each of these 270 titles was then submitted to the PsycINFO and Web of Science online databases, using both forward and backward search procedures to identify other potentially relevant articles. Specifically, the reference section of each of the 270 articles was examined for germane titles and abstracts, identifying previous studies relevant to well-being and health that were not included in the two 2005 reviews (backward search); and more recent articles were identified that cited the original 270 studies (forward search). These search procedures identified an additional 90 empirical studies to be examined for inclusion into the meta-analysis; thus, the number of empirical articles cited in these reviews or located as a result of searches using these reviews was 330.

As a final check, a database search of PsycINFO and Web of Science was conducted, combining five terms reflecting well-being (positive affect, subjective well-being, happiness, life satisfaction, and hedonic quality of life measures) and three terms reflecting health (physical health, physical well-being, and physical functioning). Only four additional studies were identified in this final search, suggesting that we successfully located most applicable published studies in the field. All potentially relevant studies published or posted through June 1, 2006, were evaluated for inclusion. Thus, the three search strategies yielded an original set of 334 empirical articles that were examined using our inclusion and exclusion criteria.

Inclusion and exclusion criteria for studies

Included studies. Potentially relevant studies were coded and included in the meta-analysis only if they met all six established criteria. To be included, a study had to (a) be written in English; (b) be an empirical study (rather than a literature review, meta-analysis, or theoretical paper); (c) include, as an independent variable, a subjective measure of well-being (e.g., positive affect, life satisfaction, happiness, optimism) or a positive mood or emotion manipulation (e.g., humorous films, imagining pleasant circumstances, etc.); (d) include, as the dependent variable, an objective measure of physical health (e.g., mortality/survival, respiratory functioning, endocrine and immune system functioning, pain tolerance, physical functioning) or illness (e.g., disease progression, heart disease, cancer, HIV symptoms); (e) state the specific sample group (e.g., cancer patients, healthy students, asthmatics); and (f) use experimental, ambulatory, and longitudinal methodology.¹ For studies that met these criteria, the effect size between well-being and health had to be either provided or computable from summary tables, descriptive statistics, or inferential statistics (*t*-statistics, *F*-ratios, odds-ratios, or Chi-square statistics). For studies that only reported multiple regression or probit analyses, *r* equivalent effect sizes (see Rosenthal & Rubin, 2003) were computed from exact *p*-values (if available) or conservative *p* cut-offs (e.g., 0.01, 0.05).

Excluded studies. As we were interested in the effects of well-being on health, studies were excluded if *only* ill-being constructs (e.g., depression, hostility, negative affect, anger) were measured or manipulated. Studies were also excluded if they (a) assessed only the contemporaneous correlation between well-being and health; (b) examined only cross-sectional mean differences between healthy and unhealthy samples; (c) measured health using only a self-report measure; or (d) examined the impact of physical health on well-being (rather than well-being on health).

In total, 120 of the 240 unique empirical studies analyzed by Lyubomirsky et al. (2005) and Pressman and Cohen (2005) were excluded based on the above criteria. A majority of these exclusions were due to the outcome being self-reported health or a health coping variable, the methodology used in the study being cross-sectional or correlational, or the study being focused on a proxy of health or a health behavior (e.g., physical exercise). Of the 94 additional empirical studies identified from our three primary search techniques, 64 studies were excluded for many of the same reasons above. As a result, 150 studies were included and coded in this meta-analysis, with nine of these studies having been uniquely considered by Lyubomirsky et al. (2005), 92 by Pressman and Cohen (2005), and 19 by both studies.

Computing effect sizes

Effect sizes. The specific type of effect size used in this meta-analysis was the correlation coefficient or r index. An r effect size was computed for all relations between well-being and health for each included study using the computer program *Comprehensive Meta-Analysis 2.0* (Borenstein, Hedges, Higgins, & Rothstein, 2005). If a study induced both positive and negative affect to compare post-induction health outcomes with participants' baseline health assessments, then effect sizes for positive affect on health and negative affect on health were computed separately. Conversely, if a study compared difference scores of a positive mood manipulation group with a negative mood manipulation group for a specified health outcome, then a single effect size was computed that estimates the size of the differential impact on health for positive moods compared to negative moods. Table I presents all studies included in the meta-analysis together with three important aggregated effect sizes. We expected well-being to relate positively to *and increase* health-related functioning, cardiovascular functioning, and immune system response. In contrast, we expected well-being to relate negatively to *and decrease* cardiovascular reactivity, endocrine response, physiological response, symptom response in chronic conditions, and disease progression. We anticipated the opposite for the effects of ill-being on these health outcomes. Thus, in Table I (and when effect sizes were entered into *Comprehensive Meta-Analysis 2.0*; Borenstein et al., 2005), positive values indicate that the relation was in the predicted direction, and negative values indicate that the relation was opposite to our predictions.

Unit of analysis. Our primary unit of analysis was the independent sample(s) within each study. Every independent sample was included and coded separately within each investigation. For example, many studies reported descriptive and inferential statistics separately for males *vs.* females, unhealthy (e.g., asthmatics) *vs.* healthy control groups, or respondents experiencing different levels of the independent variable (e.g., high *vs.* low positive affect). Further, individual effect sizes were calculated within each independent sample for all (measured or manipulated) well-being constructs and for all measured health outcomes. For example, if a study manipulated positive emotions to determine their effects on heart rate and blood pressure, we computed two separate effect sizes.

Then, for all estimates of central tendency and all tests of homogeneity, these multiple effect sizes were aggregated to derive a single effect size for each independent sample; consequently, each independent sample contributed only one r effect size for these analyses. However, coding for all possible well-being–health relations had implications for the types of moderator questions that this meta-analysis could address. First, for all of the sample-specific moderators we coded (see below), average effect sizes could be compared across all levels of the moderator – for example, effect sizes for healthy samples could be compared to effect sizes for unhealthy samples. Second, when examining the aggregate effect sizes by different health outcomes (both general and specific), because

Table I. Effect size estimates and sample characteristics for all studies.

Study	No. of samples	<i>n</i>	<i>Z</i>	Average <i>r</i>	Effect size <i>r</i> for different health outcomes			Health status of participants	Study design
					Short-term outcomes	Long-term outcomes	Disease/symptom control		
Affleck <i>et al.</i> (2000)	1	48	2.31	0.33			0.33	Medical patients	Ambulatory
Alden <i>et al.</i> (2001)	1	20	1.24	0.29	0.29			Healthy	Experimental
Apter <i>et al.</i> (1997)	1	21	0.32	0.07			0.07	Medical patients	Ambulatory
Avia and Kanfer (1980)	1	39	0.96	0.16	0.16			Healthy	Experimental
Bacon <i>et al.</i> (2004)	1	135	1.36	0.12		0.12		Medical patients	Ambulatory
Berg and Snyder (2006)	2	173	4.33	0.32	0.32			Healthy	Experimental
Berk <i>et al.</i> (1989)	1	10	0.91	0.33	0.33			Healthy	Experimental
Berk <i>et al.</i> (2001)	5	52	4.55	0.63	0.69			Healthy	Experimental
Boiten (1996)	1	32	-0.37	-0.10	-0.10	-0.10		Healthy	Experimental
Brosschot and Thayer (2003)	1	33	0.33	0.06		0.06		Healthy	Ambulatory
Brown (1993)	2	26	0.00	0.00	0.00			Healthy	Experimental
Brown <i>et al.</i> (2000)	1	335	-0.86	-0.05			-0.05	Medical patients	Longitudinal
Bruehl (1993)	2	80	2.00	0.23	0.23			Healthy	Experimental
Buchanon <i>et al.</i> (1999)	1	30	2.59	0.46	0.46			Healthy	Experimental
Carson <i>et al.</i> (1988)	1	32	1.00	0.27	0.33	0.23		Medical patients	Experimental
Carter <i>et al.</i> (2002)	1	20	0.03	0.01	0.01			Healthy	Experimental
Cassileth <i>et al.</i> (1985)	2	359	0.89	0.05			0.05	Medical patients	Longitudinal
Christie and Friedman (2004)	1	68	-0.19	-0.03	-0.04	0.00		Healthy	Experimental
Clark <i>et al.</i> (2001)	1	22	1.55	0.34	0.34			Healthy	Experimental
Codispoti <i>et al.</i> (2003).	1	10	0.53	0.20	0.20			Healthy	Experimental
Cogan <i>et al.</i> (1987)	2	36	3.26	0.53	0.53			Healthy	Experimental
Cohen <i>et al.</i> (2003)	2	334	5.27	0.28		0.28		Healthy	Experimental
Danner <i>et al.</i> (2001)	1	180	4.24	0.31		0.31		Healthy	Longitudinal
Davidson <i>et al.</i> (2003)	1	35	1.92	0.33	0.33			Healthy	Experimental
Deeg and Zonneveld (1989)	4	2,645	6.25	0.12		0.12		Mix	Longitudinal
Derogatis <i>et al.</i> (1979)	1	35	1.89	0.32			0.32	Medical patients	Longitudinal
Devins <i>et al.</i> (1990)	1	97	-0.73	-0.08		-0.08		Medical patients	Longitudinal
Dillon <i>et al.</i> (1985)	1	10	1.94	0.62	0.62			Healthy	Experimental
Ekman <i>et al.</i> (1983)	1	16	-0.84	-0.23	-0.23			Healthy	Experimental

Table I. (Continued)

Evans et al. (1993)	1	12	-0.42	-0.14	-0.14			Healthy	Ambulatory
Florin et al. (1985)	2	72	-0.89	-0.16	-0.06	-0.04	-0.44	Mixed	Experimental
Florin et al. (1985)	2	72	-0.89	-0.16	-0.06	-0.04	-0.44	Mixed	Experimental
Foster et al. (2003)	3	23	-3.97	-0.79	-0.79			Healthy	Experimental
Frazier et al. (2004)	1	56	1.37	0.19	0.19			Healthy	Experimental
Fredricksons and Levenson (1998)	2	60	3.08	0.40	0.40			Healthy	Experimental
Fredricksons et al. (2000)	6	522	1.91	0.08	0.08			Healthy	Experimental
Friedman et al. (1993)	2	1,178	-3.25	-0.09		-0.09		Healthy	Longitudinal
Futterman et al. (1992)	1	5	0.16	0.12	0.12			Healthy	Experimental
Futterman et al. (1994)	1	16	3.57	0.76	0.76			Healthy	Experimental
Gellman et al. (1990)	1	50	-0.52	-0.08	-0.08			Unhealthy community	Ambulatory
Gendolla and Krusken (2001a)	2	112	3.70	0.34	0.34			Healthy	Experimental
Gendolla and Krusken (2001b)	1	60	1.69	0.22	0.22			Healthy	Experimental
Gendolla and Krusken (2002)	2	92	0.61	0.07	0.07			Healthy	Experimental
Gendolla et al. (2001)	2	42	0.12	0.02	0.02			Healthy	Experimental
Giltay et al. (2004)	1	891	3.29	0.11		0.11		Mix	Longitudinal
Gomez (2005)	2	72	1.22	0.23	0.14	0.37		Healthy	Experimental
Gullette et al. (1997)	1	58	1.20	0.16			0.16	Medical patients	Ambulatory
Harrison et al. (2000)	1	30	0.42	0.08	0.08			Healthy	Experimental
Hertel and Hekmat (1994)	1	20	2.37	0.52	0.52			Healthy	Experimental
Hess et al. (1992)	1	27	0.40	0.08	0.08			Healthy	Experimental
Horan and Dellinger (1974)	2	24	2.00	0.44	0.44			Healthy	Experimental
Houghton et al. (2002)	3	20	0.48	0.14	0.14			Unhealthy community	Experimental
Hubert and de Jong-Meyer (1990)	1	24	0.60	0.13	0.13			Mix	Experimental
Hubert and de Jong-Meyer (1991)	1	20	1.15	0.27	0.27			Healthy	Experimental
Hubert et al. (1993)	1	52	-4.25	-0.54	-0.54			Healthy	Experimental
Hucklebride et al. (2000)	2	43	0.55	0.09	0.09			Healthy	Experimental
Hudak et al. (1991)	1	31	1.96	0.36	0.36			Healthy	Experimental
Hyland (1990)	1	10	0.53	0.20			0.20	Medical patients	Ambulatory
Jacob et al. (1999)	1	69	-1.74	-0.21	-0.21			Healthy	Ambulatory
Kawamota and Doi (2002)	1	2,274	2.72	0.06		0.06		Healthy	Longitudinal
Kitmata (2004)	4	70	2.50	0.32		0.00	0.58	Mixed	Experimental
Kivimäki et al. (2005)	2	2,852	0.53	0.01		0.01		Mix	Longitudinal
Knapp et al. (1992)	1	20	-0.35	-0.09	-0.09			Healthy	Experimental
Koivumaa-Honkanen et al. (2000)	2	7,979	16.94	0.19		0.19		Healthy	Longitudinal
Krause et al. (1997)	1	345	2.93	0.16			0.16	Medical patients	Longitudinal
Kubzansky et al. (2002)	1	455	2.22	0.10		0.10		Healthy	Longitudinal

Table I. (Continued)

Kubzansky <i>et al.</i> (2001)	1	875	6.79	0.23		0.23	Healthy	Longitudinal	
Kugler and Kalveram (1987)	1	20	1.01	0.24	0.24		Healthy	Ambulatory	
Laidlaw <i>et al.</i> (1994)	1	7	1.05	0.48		0.48	Healthy	Ambulatory	
Laidlaw <i>et al.</i> (1996)	1	38	2.63	0.42		0.42	Medical patients	Experimental	
Lambert and Lambert (1995)	1	39	2.52	0.40	0.40		Healthy	Experimental	
Lefcourt <i>et al.</i> (1990)	3	120	5.28	0.46	0.46		Healthy	Experimental	
Levenson <i>et al.</i> (1990)	1	62	0.65	0.08	0.08		Healthy	Experimental	
Levy <i>et al.</i> (1988)	1	36	3.14	0.50		0.50	Medical patients	Longitudinal	
Levy <i>et al.</i> (2002)	1	660	6.55	0.25		0.25	Mix	Longitudinal	
Liangas <i>et al.</i> (2003)	1	22	-2.69	-0.55		-0.55	Asthmatics	Ambulatory	
Lutgendorf <i>et al.</i> (1999)	1	58	2.46	0.32	0.32		Mix	Longitudinal	
Maier and Smith (1999)	1	513	6.87	0.30		0.30	Mix	Longitudinal	
McClelland and Cheriff (1997)	3	109	4.66	0.44	0.44		Healthy	Experimental	
McCraty <i>et al.</i> (1995)	1	12	-1.14	-0.36	-0.36		Healthy	Experimental	
McCraty <i>et al.</i> (1996)	1	10	1.44	0.50	0.50		Healthy	Experimental	
Meagher <i>et al.</i> (2001)	4	92	1.43	0.16	0.16		Healthy	Experimental	
Meininger <i>et al.</i> (2004)	1	371	-1.40	-0.07	-0.07		Healthy	Ambulatory	
Milam <i>et al.</i> (2004)	1	412	1.01	0.05	0.05		HIV patients	Longitudinal	
Miller and Wood (1997)	1	48	0.92	0.20	0.33		0.06	Asthmatics	Experimental
Mittwoch-Jaffe <i>et al.</i> (1995)	1	123	3.10	0.28	0.28		Healthy	Experimental	
Moskowitz (2003)	1	407	3.72	0.18		0.18	Medical patients	Longitudinal	
Neumann and Waldstein (2001)	2	42	-6.30	-0.78	-0.78		Healthy	Experimental	
Njus <i>et al.</i> (1996)	2	50	0.38	0.06	0.06		Healthy	Experimental	
O'Connor and Vallerand (1998)	1	128	1.87	0.17		0.17	Healthy	Longitudinal	
Ong and Allaire (2005)	1	33	0.53	0.10	0.10		Healthy	Ambulatory	
Ostir <i>et al.</i> (2000)	1	1,196	7.83	0.22		0.22	Healthy	Longitudinal	
Ostir <i>et al.</i> (2001)	2	2,478	3.68	0.07		0.07	Healthy	Longitudinal	
Ostir <i>et al.</i> (2004)	1	1,558	3.88	0.10		0.10	Healthy	Longitudinal	
Ostir <i>et al.</i> (2002)	1	211	3.90	0.26		0.26	Unhealthy community	Longitudinal	
Palmore (1969)	1	265	2.26	0.14		0.14	Mix	Longitudinal	
Parker <i>et al.</i> (1992)	2	421	2.32	0.11		0.11	Healthy	Longitudinal	
Perera <i>et al.</i> (1998)	1	15	2.71	0.65	0.65		Healthy	Experimental	
Pitkala <i>et al.</i> (2004)	1	491	2.87	0.13		0.13	Healthy	Longitudinal	
Polk <i>et al.</i> (2005)	1	334	2.19	0.12	0.12		Healthy	Longitudinal	
Pollard and Schwartz (2003)	1	564	0.84	0.05	0.06	0.03	Healthy	Ambulatory	
Prkachin <i>et al.</i> (1999)	1	31	1.71	0.31	0.31		Healthy	Experimental	

Table I. (Continued)

Provost and Decarie (1979)	1	40	1.44	0.23		0.23		Healthy	Experimental
Reynolds and Nelson (1981)	1	193	1.66	0.12			0.12	Medical patients	Longitudinal
Rhudy et al. (2005)	1	28	3.46	0.60	0.60			Healthy	Experimental
Richman et al. (2005)	1	1,388	5.58	0.21		0.16	0.26	Medical patients	Longitudinal
Ritz et al. (2000)	2	48	-0.93	-0.14		-0.13	-0.40	Asthmatics vs. healthy	Experimental
Ritz et al. (2001)	2	40	-0.65	-0.11			-0.11	Mix	Experimental
Ritz et al. (2005)	2	60	0.10	0.01		0.07	-0.05	Mix	Experimental
Rosenbaum (1980)	2	40	1.95	0.32	0.32			Healthy	Experimental
Santibanez-H and Bloch (1986)	1	34	-1.97	-0.34	-0.34			Healthy	Experimental
Scheier et al. (1989)	1	51	1.76	0.25			0.25	Medical patients	Longitudinal
Schwartz et al. (1981)	1	32	-0.65	-0.12	-0.12			Healthy	Experimental
Schwartz et al. (1994)	1	246	-1.19	-0.08	-0.08			Healthy	Ambulatory
Scott and Barber (1977)	2	80	1.27	0.15	0.15			Healthy	Experimental
Shapiro et al. (2001)	1	203	0.01	0.00	0.00			Healthy	Ambulatory
Sinha et al. (1992)	1	54	-2.02	-0.39	-0.36	-0.18		Healthy	Experimental
Smyth et al. (1998)	1	120	0.87	0.08	0.08			Healthy	Ambulatory
Sternbach (1962)	1	10	0.00	0.00	0.00			Healthy	Experimental
Stephoe and Holmes (1985)	2	14	-0.21	-0.07		-0.07	-0.08	Asthmatics vs. healthy	Ambulatory
Steptow and Wardle (2005)	1	160	1.74	0.14	0.14			Healthy	Longitudinal
Stevens et al. (1989)	1	20	1.21	0.29	0.29			Healthy	Experimental
Stone et al. (1994)	1	96	0.46	0.05	0.05			Mix	Ambulatory
Stone et al. (1987)	1	29	0.91	0.18	0.18			Healthy	Ambulatory
Stones et al. (1989)	1	156	-1.49	-0.12			-0.12	Institutional residents	Longitudinal
Szczepanski et al. (1997)	1	101	0.00	0.00	0.00			Healthy	Ambulatory
Uchiyama (1992)	1	6	-0.17	-0.10	-0.10			Healthy	Experimental
Uchiyama et al. (1990)	1	10	0.83	0.30	0.30			Healthy	Experimental
Van Domburg (2001)	2	354	1.71	0.09			0.09	Medical patients	Longitudinal
Van Eck et al. (1996)	1	86	0.00	0.00	0.00			Healthy	Ambulatory
Von Kanel et al. (2005)	1	27	0.00	0.00	0.00			Healthy	Experimental
Von Leupoldt and Dahme (2004)	1	20	-0.08	-0.02		-0.02		Healthy	Experimental
Von Leupoldt and Dahme (2005)	2	128	-0.92	-0.12	-0.17	0.02	-0.03	Mix	Experimental
Waldstein et al. (2000)	1	30	1.02	0.19	0.19			Healthy	Experimental
Weaver and Zillmann (1994)	2	48	1.39	0.21	0.21			Healthy	Experimental
Weid and Verbaten (2001)	1	43	1.57	0.24	0.24			Healthy	Experimental
Weisenberg et al. (1998)	1	86	2.34	0.25	0.25			Healthy	Experimental
Whorwell et al. (1992)	1	18	2.62	0.59			0.59	Medical patients	Experimental
Williams et al. (1993)	1	82	0.53	0.06		0.06		Healthy	Longitudinal

Table I. (Continued)

Wingard et al. (1994)	1	4,725	1.79	0.03		0.03	Healthy	Longitudinal
Witvliet and Vrana (1995)	1	48	0.47	0.07	0.07		Healthy	Experimental
Worthington and Shumate (1981)	4	96	3.72	0.38	0.38		Healthy	Experimental
Yogo et al. (1995)	2	24	-5.08	-0.83	-0.83		Healthy	Experimental
Yoshino (1996)	2	57	1.40	0.19	0.19		Mix	Experimental
Zachariae et al. (1991)	1	12	0.76	0.25	0.25		Healthy	Experimental
Zachariae et al. (2001)	1	15	0.40	0.11		0.11	Healthy	Experimental
Zelman et al. (1991)	1	41	2.53	0.39	0.39		Healthy	Experimental
Zillmann et al. (1993)	2	40	2.55	0.41	0.41		Healthy	Experimental
Zillmann et al. (1996)	1	43	0.79	0.12	0.12		Healthy	Experimental
Zuckerman et al. (1984)	2	351	5.20	0.27		0.25	Mix	Longitudinal
Zweyer et al. (2004)	1	56	3.73	0.47	0.47		Healthy	Experimental
Total/average	212	45,159	9.99	0.14	0.15	0.11	0.13	

Note. Each r -effect size represent the average unweighted effect size between well-being constructs and physical health outcomes within the category listed. Thus, effect sizes listed with positive values indicate enhanced health outcomes; effect sizes with negative values indicate compromised health outcomes. The sample size refers to the number of participants used to compute the effect size.

630 independent samples typically measured multiple health outcomes, we could not
statistically compare the effect sizes across different health outcomes. For
example, we could not test whether the effect of well-being on immune
635 functioning was statistically stronger or weaker than its effect on cardiovascular
functioning, because the effect sizes we used (e.g., those generated from multiple
health outcomes measured on the same sample) were not all independent from
each other. As a result, average effect sizes were not comparable across different
health outcomes, but were instead compared *within* each health outcome to
determine whether the relation between well-being and health was significantly
stronger than zero.

640 *Coding sample moderators*

All independent samples that met the inclusion criteria were coded for possible
moderators. Specifically, two classes of moderators were coded: (a) variable
characteristics (health and/or illness; state or trait well-being); and (b) sample
characteristics (health status, age, gender).

645 *Objective health outcome characteristics.* As the third goal of our meta-analysis was
to compare effect sizes for the link between well-being and health across different
health outcomes, we coded three general categories of health and 12 specific
health outcome variables. First, for short-term outcomes (health outcomes
measured at the molecular level), we coded the following: immune system
650 response (e.g., sIgA concentration, NKCA), endocrine system response (e.g.,
cortisol, epinephrine, norepinephrine), cardiovascular system reactivity (e.g.,
blood pressure, heart rate), physiological response (e.g., finger temperature, skin
conductance), and pain tolerance (e.g., time in cold pressure task). Second, for
long-term outcomes (health outcomes measured at the molar level, for normal
655 functioning) we coded the following: general health (e.g., accumulated mucus
weight during infections, cholesterol), cardiovascular system functioning (e.g.,
high and low frequency power, ischemic episodes), respiratory functioning (e.g.,
inspiration volume, inspiratory volume), and longevity (mortality for participants
without a chronic condition). Third, for chronic conditions (health outcomes
660 measured at the molar level, for those with chronic illness), we coded respiratory
diseases/conditions (e.g., flare reactions, wheal reactions, forced expiratory
volume, bronchial responsiveness), disease progression (e.g., complete recovery
from disease, viral load), and survival (staying alive despite having a chronic
condition).

665 *State and trait measures of well-being.* We recorded the exact SWB construct
measured or manipulated in each study. Each of these constructs was then coded
as representing either a state variable (momentary positive and negative affect,
induced hope, relaxation, etc.) or a trait variable (global life satisfaction,
happiness, optimism, and trait levels of positive and negative affect) of well-
670 being or ill-being.

Sample characteristics. We predicted that the effects of well-being on objective health outcomes would be moderated by characteristics of the independent samples. Two sample-level categorical moderators were coded: (a) method of study (experiment, ambulatory, or longitudinal); and (b) health status of the sample at baseline (healthy or unhealthy). In addition, we recorded two continuous sample-level moderators: (a) mean participant age at baseline; and (b) gender composition (percent male respondents). Coding of the sample characteristics was based on information gleaned from the Method section of each article.

Data analysis: Fixed vs. random effects

Both fixed and random effects methods for meta-analysis have advantages. The fixed effects model provides a more precise and reliable estimate of the population effect size (Cooper, 1998), whereas the random effects model allows for relatively more generalizable conclusions. Random effects models also specify the amount of variance accounted for by between-studies differences and variance accounted for by within-study differences. As each has advantages and disadvantages, aggregate r effect sizes were computed with *Comprehensive Meta-Analysis 2.0* (Borenstein et al., 2005) using both fixed and random models. When possible, we focused on the results from the random effect models. However, some moderator tests (e.g., meta-regressions) can only be estimated using fixed effects models. In these cases, all null hypotheses, fixed effects models, and *post hoc* comparisons followed steps outlined by Hedges (1994). For both models, homogeneity tests were used to determine whether variance in the effect sizes was explained by the proposed moderators. When the r effect sizes were aggregated using a fixed effects model, measures of central tendency were calculated by averaging weighted r effect sizes (inverse variance weights) across all independent samples. When categorical groups were independent, and if a categorical moderator explained significant variance in the effect sizes (i.e., $p < 0.05$ for Q_{BET}), then *post hoc* contrasts were performed to determine which groups were statistically different. For continuous moderators, meta-regression analyses were used to determine whether variation in the effect sizes was explained by the moderator.

Results

Description of the literature included

Publication statistics. Our search techniques identified a total of 150 studies that met the established inclusion criteria. From these studies, 212 independent samples were identified, from 17 (mostly Western) nations, which measured or manipulated a well-being construct and measured a physical health outcome. From these 212 samples, 439 distinct effect sizes were computed between well-being and physical health; an additional 310 effect sizes were computed between ill-being and health from the 79 studies that measured both well-being and ill-being. The number of independent samples coded per study ranged from 1 to 6,

with 107 studies (71.3%) using a single independent sample, and 33 studies (22.0%) reporting two independent samples (see Table I for a detailed description of each study). The number of effect sizes of the link between well-being and physical health per study ranged from 1 to 20 ($M=2.93$, $SD=2.44$), with 70.7% of the studies reporting 3 or fewer effect sizes. The typical study surveyed 50 respondents ($M=50.50$); however, the data with respect to sample size were positively skewed ($M=294$, $SD=867$).

Characteristics of the independent samples. The 150 studies recruited a total of 44,159 respondents. Some 114 studies ($N=35,863$) reported participant age ($M=37.91$ years, $SD=20.23$). Of the 143 studies that reported gender composition, 30.8% had nearly equal numbers of males and females (45–55% male), with the majority of the remainder (61.6%) reporting a higher percentage of females (at least 56%). Table II summarizes in detail several other sample characteristics. For example, a majority of the studies were published during the last 15 years (76.0%), employed an experimental design (59.3%), included either a student or community sample (82.7%), received funding from academic or government sources (60.7%), and were conducted in the United States (56.7%).

Meta-analyzing the samples

What is the overall effect of well-being on health? As shown in Table III, from the 212 independent samples, the mean unweighted r effect size for the well-being–health relation was 0.135 (95% CI=0.110–0.160) from the random effects analysis, and 0.115 from the fixed effects analysis. Both r effect sizes are significantly different from zero ($Z=9.99$ and 23.69, respectively). In addition, the second goal of this meta-analysis was to compare the effect of well-being on health to that of ill-being on health. Using the 99 independent samples that measured the impact of both well-being *and* ill-being on health, we found the ill-being effect size to be significant, negative, and of approximately the same magnitude ($r_{random} = -0.155$; $r_{fixed} = -0.099$, both $ps < 0.001$) as the well-being effect.

Notably, the average well-being–health effect sizes were not consistent across all sample characteristics (see Q -values in Table III). Omnibus homogeneity tests demonstrated substantial within-group variation across the 212 independent samples that measured well-being ($Q_w [211] = 903.88$, $p < 0.001$). Hence, we examined average effect sizes separately for each of the three study designs (experimental, ambulatory, and longitudinal). Even with the less powerful random effects model, the average effect sizes varied by study design ($Q_{BET} [2, k=212] = 10.50$, $p = 0.005$). Studies that used ambulatory procedures ($r_{random} = 0.029$, ns) reported significantly lower effect sizes than both longitudinal and experimental studies. With the fixed effects model, the average effect of well-being on health was smaller for longitudinal designs ($r_{fixed} = 0.113$) than for experimental designs ($r_{fixed} = 0.166$). As study design proved to be a moderator of the well-being–health effect sizes, and experimental procedures provide the only

Table II. General characteristics of included studies.

Characteristic	No. of studies (%)	<i>r</i> Effect size	Total <i>n</i>
Year of report			
2001–2006	53 (35.3)	0.11	19,173
1991–2000	61 (40.7)	0.15	15,143
1981–1990	28 (18.7)	0.15	9,310
1970–1980	6 (4.0)	0.27	258
Before 1970	2 (1.3)	0.07	275
Design of study			
Experimental	89 (59.3)	0.16	4,683
Longitudinal	38 (25.3)	0.14	37,128
Ambulatory	23 (15.3)	0.04	2,348
Population sampled			
Community	65 (43.3)	0.11	37,158
Students	59 (39.3)	0.15	3,162
Children/adolescents	6 (4.0)	−0.05	1,730
Mixed/specialized	20 (13.3)	0.25	2,109
Funding source			
None	59 (39.3)	0.17	9,454
NIMH	9 (6.0)	0.18	1,620
NIA	8 (5.3)	0.18	8,183
NIH	6 (4.0)	0.09	5,713
Academic institution	6 (4.0)	−0.10	276
Grant – other	55 (36.7)	0.14	18,050
Most common journals			
<i>Psychosomatic Medicine</i>	20 (13.3)	0.12	4,891
<i>Psychophysiology</i>	10 (6.7)	0.11	727
<i>Biological Psychology</i>	5 (3.3)	0.14	199
<i>Journal of Psychosomatic Research</i>	5 (3.3)	0.02	444
Site of study			
United States	85 (56.7)	0.16	22,165
Germany	14 (9.3)	0.03	1,249
England	12 (8.0)	0.16	956
Netherlands	7 (4.7)	0.07	4,084
Canada	6 (4.0)	0.16	970
Japan	6 (4.0)	−0.01	2,441

Note. Each *r*-effect size represents the average unweighted effect size between well-being constructs and physical health outcomes within the category listed.

direct test of causal pathways, many of the subsequent analyses were performed with the three study designs combined and with the experimental studies alone.

Does well-being differentially impact health outcomes? The second goal of this meta-analysis was to determine which health outcomes were most strongly associated with well-being. First, we examined how well-being (and ill-being) was related to our general health categories (short-term outcomes, long-term outcomes, and disease/symptom control). As shown in Table IV, an analysis with 141 samples demonstrated that increases in well-being were positively associated with short-term outcomes ($r_{\text{random}} = 0.148$; $p < 0.001$). Notably, this effect was slightly stronger for the 123 studies that used experimental designs ($r_{\text{random}} = 0.172$,

Table III. The effect of well-being and ill-being on health outcomes by study design.

Study design	Sample size		<i>r</i> Effect size ^a		95% CI random effects model		Z-value		Test of heterogeneity	
	<i>n</i>	<i>K</i>	Random	Fixed	Lower	Upper	Random	Fixed	<i>Q</i> -value	Significance
Well-being	42,928	212	0.135	0.115	0.110	0.160	9.99	23.688	903.88	<0.001
Experimental	4,428	139	0.164 _A	0.166 _A	0.126	0.202	8.366	11.162	442.475	<0.001
Ambulatory	2,066	24	0.029 _B	-0.005 _C	-0.035	0.102	0.768	-0.256	47.184	0.012
Longitudinal	36,434	49	0.128 _A	0.113 _B	0.090	0.166	6.556	21.810	356.861	<0.001
Ill-being	8,187	99	-0.155	-0.099	-0.113	-0.196	7.166	11.171	341.214	<0.001
Experimental	1,892	68	-0.166 _A	-0.159 _A	-0.107	-0.224	5.462	7.730	216.091	<0.001
Ambulatory	1,707	18	-0.152 _A	-0.098 _B	-0.064	-0.238	3.368	7.277	54.315	<0.001
Longitudinal	4,588	13	-0.133 _A	-0.071 _B	-0.044	-0.221	2.915	4.954	58.348	<0.001

Note. Well-being includes life satisfaction, happiness, and positive emotions, whereas ill-being comprises such negative constructs as stress, depression, and anger. Effect sizes with different subscripts in each column differed significantly at $p < 0.05$. Within the well-being and ill-being sections, effect sizes are independent across study design, so experimental, longitudinal, and ambulatory effect sizes can be compared. Across the well-being and ill-being sections, effect sizes are not independent, so comparisons cannot be made (e.g., experimental to experimental). All effect sizes with Z -values > 1.96 are significant at $p < 0.05$.

^aEffect sizes listed with positive values indicate enhanced health outcomes; effect sizes with negative values indicate compromised health outcomes.

Table IV. The effect of well-being and ill-being on three types of health outcomes.

General categories of physical health outcomes	Sample size		<i>r</i> Effect size ^a		95% CI random effects model		Z-value		Test of heterogeneity	
	<i>n</i>	<i>K</i>	Random	Fixed	Lower	Upper	Random	Fixed	<i>Q</i> -value	Significance
Well-being										
Short-term outcomes	6,430	141	0.148	0.084	0.099	0.197	5.830	7.159	489.937	<0.001
Long-term outcomes	34,106	51	0.112	0.119	0.087	0.152	7.084	21.017	337.026	<0.001
Disease/symptom control	3,623	33	0.127	0.140	0.061	0.192	3.748	8.849	99.206	<0.001
Ill-being										
Short-term outcomes	3,584	73	-0.166	-0.114	-0.105	-0.225	5.312	9.671	285.847	<0.001
Long-term outcomes	3,564	18	-0.081	-0.054	-0.018	-0.144	2.515	3.522	43.321	<0.001
Disease/symptom control	1,275	18	-0.180	-0.154	-0.082	-0.274	3.568	6.220	51.643	<0.001

Note. Effect sizes within each category are not independent and cannot be compared. All effect sizes with *Z*-values >1.96 are significant at $p < 0.05$.

^aEffect sizes listed with positive values indicate enhanced health outcomes; effect sizes with negative values indicate compromised health outcomes.

$p < 0.001$). Also, the associations between well-being and long-term outcomes ($r_{\text{random}} = 0.112$; $p < 0.001$) and between well-being and disease/symptom control ($r_{\text{random}} = 0.127$; $p < 0.001$) were both positive, suggesting that well-being promoted healthy functioning and symptom control. However, when examining these associations with only experimental studies, the well-being – long-term outcomes average effect size was smaller and non-significant ($K = 15$; $r_{\text{random}} = 0.089$, $p = 0.11$), and the well – being-disease/symptom control average effect was nearly identical to that for the non-experimental studies but non-significant ($K = 12$; $r_{\text{random}} = 0.122$, $p = 0.21$).

Does ill-being differentially impact health outcomes? As expected, ill-being was negatively related to each category of health outcomes (see Table IV). Comparisons of the effects of ill-being vs. well-being revealed that ill-being has a slightly stronger effect on short-term outcomes ($r_{\text{random}} = -0.166$ vs. $r_{\text{random}} = 0.148$) and disease/symptom control ($r_{\text{random}} = -0.180$ vs. $r_{\text{random}} = 0.127$), whereas well-being has a slightly stronger effect on long-term outcomes ($r_{\text{random}} = 0.112$ vs. $r_{\text{random}} = -0.081$). Thus, in general, the effect sizes for both well-being and ill-being were rather similar (though in opposite directions). These relations held for the experimental studies. Specifically, the average effect size for the short-term outcomes assessed from the 58 samples that manipulated ill-being ($r_{\text{random}} = -0.171$) was nearly identical to the effect size from the 123 samples that manipulated well-being ($r_{\text{random}} = 0.172$). This finding demonstrates that inductions of well-being lead to healthy functioning and inductions of ill-being lead to compromised health at similar magnitudes.

Does well-being differentially impact specific types of health outcomes? The third goal of this meta-analysis was to determine which types of health outcomes were most strongly associated with well-being. As displayed in Table V, we examined the effects of well-being on 12 specific health outcomes. Focusing on the random effects models for the five short-term health outcomes, we observe that the specific health outcome explained a significant amount of the heterogeneity ($Q_{\text{BET}} [5, k = 141] = 131.509$, $p < 0.001$). Well-being was strongly associated with improved immune functioning ($r_{\text{random}} = 0.332$) and higher pain tolerance ($r_{\text{random}} = 0.320$). As expected, well-being was also associated with a decreased endocrine system response ($r_{\text{random}} = -0.101$), although this relation was much weaker and only marginally significant when compared to immune and pain outcomes. Finally, well-being was not associated with cardiovascular response/reactivity ($r_{\text{random}} = 0.026$) nor physiological response ($r_{\text{random}} = -0.031$).

Average effect sizes were more homogenous for long-term outcomes and disease/symptom control. For long-term outcomes, well-being was most strongly associated with increased longevity ($r_{\text{random}} = 0.137$). Well-being also predicted improved general health ($r_{\text{random}} = 0.110$) and cardiovascular functioning ($r_{\text{random}} = 0.119$), and was marginally related to better respiratory functioning ($r_{\text{random}} = 0.071$; $p < 0.10$). For disease/symptom control, well-being was associated with slower disease progression ($r_{\text{random}} = -0.150$) and longer survival

Table V. The effect of well-being and ill-being on specific health outcomes.

Specific measures of physical health outcomes	Sample size		<i>r</i> Effect size ^a		95% CI random effects model		Z-value ^b		Test of heterogeneity	
	<i>n</i>	<i>K</i>	Random	Fixed	Lower	Upper	Random	Fixed	<i>Q</i> -value	Significance
Short-term outcomes										
Immune system response	1,323	32	0.332	0.224	0.228	0.410	6.423	8.110	73.529	<0.001
Pain tolerance	1,096	37	0.320	0.320	0.257	0.380	9.467	10.505	41.504	0.243
Endocrine system response ^c	1,154	21	-0.101	-0.090	0.001	-0.201	1.939	2.968	40.607	0.004
Cardiovascular system reactivity ^d	3,181	60	0.026	0.018	-0.045	0.096	-0.710	-1.144	221.769	<0.001
Physiological response	527	18	-0.031	-0.056	0.098	-0.156	0.473	1.343	36.596	0.004
Long-term outcomes										
Cardiovascular functioning	4,332	10	0.119	0.117	0.056	0.181	3.706	7.885	25.445	0.003
General health	5,124	7	0.110	0.057	0.024	0.195	2.511	4.110	33.072	<0.001
Longevity ^e	24,869	24	0.137	0.128	0.093	0.181	5.989	20.435	263.246	<0.001
Respiratory functioning	672	12	0.071	0.071	-0.002	0.144	1.907	1.907	7.721	0.738
Disease/symptom control										
Respiratory conditions	353	16	-0.105	-0.129	0.056	-0.262	1.281	2.841	39.671	0.001
Disease progression	1,540	8	-0.150	-0.170	-0.018	-0.276	2.229	6.908	36.137	<0.001
Survival ^e	2,065	10	0.097	0.093	0.018	0.175	2.394	4.420	28.330	0.001

Note. Effect sizes from the categories of health are not independent and cannot be compared. All effect sizes with *Z*-values >1.96 are significant at *p* <0.05.

^aPositive values indicate that well-being produces increased levels of the health category; negative values indicate that well-being produces decreased levels of the health category. ^bPositive *Z*-values indicate that results were in the expected directions. For example, we would expect well-being to produce less cardiovascular reactivity, but there is a non-significant increase, so the *Z* is a negative value. ^cRefers to stress hormones, such as cortisol and epinephrine. ^dIncludes heart rate reactivity and blood pressure responses. ^eLongevity refers to overall length of life. Survival refers to staying alive despite having one or more chronic conditions.

Well-being and health: A meta-analysis 29

from chronic illness ($r_{random} = 0.097$). Although well-being was not significantly related to reduced respiratory conditions ($p = 0.20$ from the random effects model), the effect size was in the predicted direction ($r_{random} = -0.105$).

All of the above results – for the short-term, long-term, and disease control categories – were minimally altered when the ambulatory and longitudinal studies were removed, with the exceptions that the positive impact of well-being on improved immune functioning ($r_{random} = 0.371$) and better general health ($r_{random} = 0.283$) both became stronger, while the positive impact of well-being on long-term cardiovascular functioning ($r_{random} = 0.016$) and long-term respiratory functioning became weaker ($r_{random} = 0.018$) and non-significant.

Sample specific moderators of the well-being–health associations

As demonstrated by the Q -values in Tables IV and V, tests of heterogeneity indicated that some of the aggregate effect sizes may be moderated by sample level characteristics, such as health status of the sample at baseline, average age of the respondents, and exact health outcome measured. For example, although well-being had a positive effect on all three types of health outcomes (see Table IV), significant heterogeneity was observed within each group. A closer examination by specific health outcome (see Table V) indicates that heterogeneity is minimal for some outcomes (such as pain tolerance) and much more significant for other outcomes (such as immune function and cardiovascular reactivity). Thus, based on both our general and specific categories and the heterogeneity statistics, we defined five specific groups and examined potential moderators: (a) short-term immune system functioning; (b) short-term endocrine response; (c) cardiovascular reactivity and physiological response; (d) long-term promotion of healthy functioning (including cardiovascular functioning, general health, and mortality); and (e) enhanced symptom control and survival in chronic conditions.

To explore what underlies this heterogeneity, we focused on five moderators (three categorical and two continuous). The categorical moderators were (a) health status of the sample at baseline (healthy or unhealthy), (b) exact type of health outcome measured (e.g., sIgA antibody production in immune system response; heart rate and blood pressure in cardiovascular reactivity), and (c) the operational definition of well-being (as a state or trait variable). To illustrate the moderating effect of these categorical variables, we reported the descriptive and inferential statistics for each level of the specified moderator. Although we reported both random and fixed effects, we focused here on the more generalizable random effect models. The continuous moderator variables were (a) average age of the sample, and (b) percent male respondents. Continuous moderator effects were examined using meta-regression analyses (a necessarily fixed effects models), focusing on the slope (β_1) of the meta-regression line, which indicates whether the effect sizes were associated with changes in the continuous moderators. In each case, we only tested for moderators if the outcome was reported in 10 or more samples. For the exact health outcomes,

when effect sizes could not be grouped from 10 independent studies, we documented the exact health outcome most commonly reported.

Moderators of well-being and short-term immune system functioning. Several characteristics of the study sample moderated the effect of well-being on short-term immune system functioning. First, although the average effect sizes for healthy and unhealthy samples were both positive (e.g., increases in well-being were associated with improved immunity), the average effect size was significant for healthy ($r_{random} = 0.360$) but not unhealthy ($r_{random} = 0.147$) samples (see top section of Table VI). Further, this positive effect of well-being was magnified in studies measuring sIgA antibody production. For these studies, the average impact of well-being on short-term immune system functioning was stronger ($r_{random} = 0.370$) than that in studies that measured other markers of normal immune responses (e.g., increased t cell counts on markers such as CD4, CD8+, and CD16+; $r_{random} = 0.257$). Additionally, studies that manipulated or measured state well-being variables reported higher average effect sizes ($r_{random} = 0.338$) than studies that determined the relation between well-being and immune system functioning using trait measures of well-being ($r_{random} = 0.164$, *ns*). Finally, the gender composition of the sample also moderated the relation between well-being and short-term immune system functioning (see top section of Table VII). The slope of the meta-regression line is significantly negative ($\beta_1 = -0.267$, $p = 0.0028$), which indicates that samples with a higher proportion of female respondents reported larger effect sizes on average.²

Moderators of well-being and short-term endocrine response. Several characteristics of the study sample also moderated the effect of well-being on short-term endocrine response (see the second section of Tables VI and VII). It should be noted again that a negative relation between well-being and endocrine response should be interpreted as promoting healthy functioning, because the increase in endocrine response from negative affect is typically interpreted as compromising health (especially when stress hormones are measured). With respect to the categorical variables, the effect of well-being on decreased endocrine response was not significant in the healthy sample group ($r_{random} = -0.075$, $p = 0.16$). The effect size for the unhealthy sample group was quite a bit larger ($r_{random} = -0.343$); however, the unhealthy sample group consisted of only a single small study ($N = 26$). Thus, neither of these effect sizes was significant. As was observed in the relation between well-being and immune system functioning, the negative effect of well-being was strongest (and significant) when studies assessed the most commonly measured hormone associated with stress – levels of cortisol. For these studies, the average impact of well-being on short-term endocrine response was stronger ($r_{random} = -0.109$, $p = 0.04$) than that for studies that measured other stress hormones ($r_{random} = -0.043$, *ns*). Finally, neither the sample's gender composition nor the average age of the respondent moderated the relation between well-being and short-term endocrine response (see the

Table VI. Categorical moderators of relations between well-being and grouped health outcomes.

	Sample size		<i>r</i> Effect size ^a		95% CI random effects model		Z-value ^b		Test of heterogeneity	
	<i>n</i>	<i>K</i>	Random	Fixed	Lower	Upper	Random	Fixed	<i>Q</i> -value	Significance
Well-being-short – term immune system functioning										
Moderator										
Health status										
Sample healthy	715	27	0.360	0.335	0.262	0.451	6.795	9.048	46.483	0.008
Sample unhealthy	454	3	0.147	0.070	-0.117	0.393	1.090	1.470	2.527	0.283
Exact health outcome										
SigA	514	16	0.370	0.327	0.252	0.478	5.792	7.646	29.439	0.014
All other	853	18	0.257	0.153	0.113	0.391	3.435	4.323	40.319	0.001
State or trait SWB										
State	853	30	0.338	0.301	0.243	0.427	6.612	8.82	54.439	0.003
Trait	470	2	0.164	0.083	-0.119	0.422	1.139	1.797	3.844	0.050
Well-being-short-term endocrine response										
Health status										
Sample healthy	19	1,104	-0.075	-0.077	0.030	-0.178	1.398	2.493	36.273	0.007
Sample unhealthy	1	26	-0.343	-0.343	0.138	-0.693	1.449	1.716	-	-
Exact health outcome										
Cortisol	21	1,154	-0.109	-0.092	-0.003	-0.212	2.020	3.044	43.488	0.002
All other	4	133	-0.043	-0.043	0.135	-0.217	0.471	0.471	0.952	0.813
State or trait SWB										
State	19	660	-0.097	-0.059	0.022	-0.214	1.593	1.447	39.257	0.003
Trait	2	494	-0.131	-0.127	0.114	-0.361	1.048	2.830	0.058	0.809
Well-being-cardiovascular reactivity and physiological response										
Moderator										
Health status										
Sample healthy	3,124	57	0.014	0.011	-0.058	0.085	-0.373	-0.731	215.567	<0.001
Sample unhealthy	140	5	-0.011	0.013	0.252	-0.272	0.081	-0.145	8.575	0.073
Exact health outcome										
Blood pressure	2,218	32	0.091	0.064	-0.005	0.186	-1.868	-3.597	156.792	<0.001
Heart rate	1,841	43	0.060	0.019	-0.034	0.154	-1.251	-0.884	155.314	<0.001
Skin conductance	396	16	0.016	-0.016	-0.114	0.145	-0.236	0.368	30.094	0.012
State or trait SWB										
State	3,128	62	0.016	0.017	-0.054	0.086	-0.449	-1.100	72.342	<0.001
Trait	160	1	-0.136	-0.136	0.019	-0.285	1.717	1.717	-	-

Table VI. (Continued)

	Sample size		<i>r</i> Effect size ^a		95% CI random effects model		<i>Z</i> -value ^b		Test of heterogeneity	
	<i>n</i>	<i>K</i>	Random	Fixed	Lower	Upper	Random	Fixed	<i>Q</i> -value	Significance
Well-being – long-term healthy functioning										
Health status										
Sample healthy	23	24,315	0.113	0.114	0.066	0.160	4.677	17.876	236.784	<0.001
Sample unhealthy	4	942	0.086	0.112	−0.032	0.203	1.428	3.836	8.387	0.039
Exact health outcome										
Coronary risk factors ^c	7	4,480	−0.125	−0.059	−0.134	0.190	3.693	8.224	17.113	0.009
Other	10	4,897	0.114	0.063	0.049	0.179	3.397	5.048	39.876	<0.001
State or trait SWB										
State	7	567	0.075	0.075	0.002	0.147	2.014	2.043	6.077	0.415
Trait	32	32,867	0.132	0.114	0.095	0.168	6.946	20.888	320.828	<0.001
Well-being – enhanced symptom control and survival during chronic conditions										
Moderator										
Health status										
Sample healthy	42	3	0.095	0.078	−0.266	0.432	0.612	0.633	2.538	0.281
Sample unhealthy	3,543	29	0.120	0.138	0.052	0.187	3.445	8.633	93.261	<0.001
Exact health outcome										
Asthma symptoms	380	16	−0.077	−0.125	0.114	−0.262	0.787	2.725	52.998	0.003
Recovery from disease	1,919	10	0.145	0.126	0.063	0.224	3.463	5.520	24.925	0.003
State or trait SWB										
State	447	19	0.109	0.131	−0.041	0.225	1.424	3.118	49.760	<0.001
Trait	3,176	14	0.134	0.142	0.064	0.203	3.721	8.285	49.388	<0.001

Note. The following samples were not included in the analysis of healthy *vs.* unhealthy samples because they combined healthy and unhealthy samples: two samples that measured immune functioning, one sample that measured endocrine response, 12 samples that measured long-term optimal and healthy functioning, and one sample that measured enhanced symptom control. Effect sizes from the categories of health are not independent and cannot be compared. All effect sizes with *Z*-values >1.96 are significant at *p* <0.05.

^aPositive values indicate that well-being produces higher levels of the health category; negative values indicate that well-being produces lower levels of the health category. ^bPositive *Z*-values indicate that results were in the expected directions. ^cWe coded cholesterol ratio, HDL and LDL cholesterol level, hypertension, high and low frequency power, non-fatal MI, triglycerides levels as coronary risk factors for the purposes of this moderator test.

Well-being and health: A meta-analysis 33

Table VII. Continuous moderators of relations between well-being and grouped health outcomes.

Parameter	Estimate	SE	Z-value	95% CI for β	
				LL	UL
Well-being – short-term immune system functioning					
Moderator: average age of the sample (age centered on sample mean)					
$Q_{\text{Model}}(1, k=23) = 0.738, p = 0.390$					
β_0	0.199	0.034	5.774	0.131	0.267
β_1	-0.002	0.002	-0.859	-0.006	0.002
Moderator: percentage of male respondents					
$Q_{\text{Model}}(1, k=32) = 8.91, p = 0.003$					
β_0	0.392	0.089	6.345	0.271	0.513
β_1	-0.267	0.062	-2.986	-0.092	-0.441
Well-being – short-term endocrine response					
Moderator: average age of the sample (age centered on sample mean)					
$Q_{\text{Model}}(1, k=18) = 0.228, p = 0.633$					
β_0	-0.073	0.033	2.152	-0.006	-0.139
β_1	-0.001	0.003	0.478	0.004	-0.007
Moderator: percentage of male respondents					
$Q_{\text{Model}}(1, k=20) = 0.556, p = 0.454$					
β_0	-0.118	0.056	2.104	-0.007	-0.229
β_1	0.073	0.097	0.748	-0.118	0.264
Well-being – cardiovascular reactivity and physiological response					
Moderator: average age of the sample (age centered on sample mean)					
$Q_{\text{Model}}(1, k=39) = 10.093, p = 0.001$					
β_0	0.025	0.019	-1.300	-0.012	0.062
β_1	-0.005	0.001	3.176	-0.002	-0.007
Moderator: percentage of male respondents					
$Q_{\text{Model}}(1, k=61) = 3.315, p = 0.068$					
β_0	-0.050	0.037	1.376	0.021	-0.122
β_1	0.122	0.067	-1.82	-0.009	0.251
Well-being – long-term promotion of healthy functioning					
Moderator: average age of the sample (age centered on sample mean)					
$Q_{\text{Model}}(1, k=35) = 0.450, p = 0.502$					
β_0	0.128	0.006	21.721	0.116	0.140
β_1	0.0002	0.0003	0.671	-0.0004	0.0008
Moderator: percentage of male respondents					
$Q_{\text{Model}}(1, k=36) = 21.187, p < 0.001$					
β_0	0.089	0.009	9.524	0.071	0.107
β_1	0.073	0.015	4.603	0.042	0.104
Well-being – enhanced symptom control and survival during chronic conditions					
Moderator: average age of the sample (age centered on sample mean)					
$Q_{\text{Model}}(1, k=28) = 2.608, p = 0.106$					
β_0	0.111	0.021	5.122	0.069	0.154
β_1	0.002	0.001	1.615	-0.0004	0.0042
Moderator: percentage of male respondents					
$Q_{\text{Model}}(1, k=32) = 0.803, p = 0.370$					
β_0	0.175	0.037	4.688	0.101	0.248
β_1	-0.048	0.053	-0.896	0.057	-0.153

Note. Z-value tests the null hypothesis that the parameter is zero in the population. Moderators that included fewer than 10 samples were not examined using the meta-regression analyses.

second section of Table VII). This indicates that the marginally significant drop in stress hormones as a result of well-being is constant across gender and age.

Moderators of well-being and short-term cardiovascular and physiological reactivity. Fewer characteristics of the study sample moderated the effect of well-being on cardiovascular and physiological reactivity (see the third section of Tables VI and VII). For example, the null association was consistent regardless of whether the sample (a) was healthy or unhealthy, (b) measured heart rate or skin conductance, or (c) manipulated state well-being.

These analyses show only one significant moderating characteristic of the sample – as the average age of the respondents increases, well-being is associated with decreases in cardiovascular reactivity and physiological response. Again, as was true of endocrine system response, a negative relation between well-being and cardiovascular reactivity and physiological response is interpreted as promoting healthy functioning. Thus, as the average age of the sample increased, well-being was associated with promoting healthy cardiovascular reactivity and physiological response. However, three marginally significant findings are of interest. First, studies that measured blood pressure in response to increases in well-being demonstrated that well-being was associated with a marginally significant *increase* in blood pressure. Second, one study measured trait levels of well-being in a follow-up analysis and found these to predict lower levels of cardiovascular reactivity and physiological response. Third, the marginally significant positive slope from the meta-regression examining gender composition ($\beta_1 = 0.122$, $p = 0.07$) indicated that samples with more females reported stronger *negative* relations between well-being and cardiovascular reactivity and physiological response.

Moderators of well-being and long-term healthy functioning. Several characteristics of the study sample moderated the effect of well-being on long-term healthy functioning. First, although the average effect sizes for both healthy and unhealthy samples were both positive (e.g., increases in well-being were associated with long-term healthy functioning), the average effect size was significant for healthy ($r_{random} = 0.113$) but not unhealthy ($r_{random} = 0.086$) samples (see Table VI). Interestingly, the different types of long-term health outcomes did not differ widely. For example, coronary risk factors ($r_{random} = -0.125$), longevity ($r_{random} = 0.137$), and other general health outcomes ($r_{random} = 0.114$) all had similar average effect sizes. Additionally, studies that measured state well-being reported lower average effect sizes ($r_{random} = 0.075$) than studies that examined the relation between well-being and long-term optimal and healthy functioning using trait measures of well-being ($r_{random} = 0.132$). Finally, although average age of the respondents was not a significant moderator, the gender composition of a sample did moderate this relation (see the fourth section of Table VII). The slope of the meta-regression line is significantly positive ($\beta_1 = 0.073$, $p < 0.001$), which suggests that the

average effect size for a sample with all females was lower ($r_{random} = 0.089$) than the average effect size for a sample with all males ($r_{random} = 0.162$).

Moderators of well-being and symptom control during chronic conditions. Several characteristics of the study sample moderated the effects of well-being on symptom control during chronic conditions. First, three small studies ($N = 20$) measured symptoms of chronic conditions (asthma and allergy symptoms) on healthy samples (see Table VI) in comparison to unhealthy samples. These studies reported effect sizes that were on average lower ($r_{random} = 0.095$) than those samples that measured chronic condition on exclusively unhealthy samples ($r_{random} = 0.120$). Also, the type of health outcome measured significantly affected the average effect size. For example, well-being had a stronger relation with recovery from disease ($r_{random} = 0.145$) than with the reduction of asthma symptoms ($r_{random} = -0.077$, $p = 0.43$). Additionally, studies that measured state well-being revealed a lower, non-significant average effect size ($r_{random} = 0.109$) than studies that measured trait well-being ($r_{random} = 0.134$). Finally, neither the samples' gender composition nor average age was found to moderate this relation (see the last section of Table VII). This indicates that the significant increase in symptom control and survival that is linked to well-being is constant across gender and age.

Discussion

This meta-analysis examined the unidirectional effect of well-being on objective physical and physiological health outcomes. Pooling the results of 150 experimental, ambulatory, and longitudinal studies, we found an average overall r effect size of 0.14 between well-being and objective health. The aggregated r -effect size for the 123 experimental studies that induced positive emotion was 0.17. Effect sizes can best be conceptualized in practical terms using a binomial effect size display (BESD; see Rosenthal, 1991, 1994, for a full explanation of the BESD procedure and rationale). The BESD is most easily understood when the outcome is dichotomous, as it is for survival. As a case in point, consider the interpretation of the aggregate r effect size between well-being and longevity ($r_{random} = 0.14$; see Table V). Using the BESD to interpret this r effect size (see Table VIII), we expect the probability of living longer increases by 14% for individuals with high well-being compared to those with low well-being. Also, we

Table VIII. Binomial effect size display for the average impact of well-being on longevity.

Levels of variable	High well-being	Low well-being
Survival	57	43
Death	43	57

Note. The BESD is based on the average effect size ($r = 0.14$) for well-being and survival (see Table V).

find that the survival rate increases 10% for individuals with a chronic illness who have high versus low well-being (BESD not pictured).

Furthermore, not only can r effect sizes be converted into a BESD, but the BESD itself can then be translated into two other effect sizes often reported in biomedical research – namely, relative risk (RR) and odds ratio (OR; see Rosenthal & DiMatteo, 2001, for the steps to convert BESD information into other effect size indices). Using the BESD presented in Table VIII, our overall effect size can be interpreted as the odds of survival (OR = 1.75) and as the relative risk of mortality (RR = 0.75). In each case, survival is more likely in the high well-being group and mortality is more likely in the low well-being group. When balancing the costs of improving well-being against the benefits of saving lives, these are very significant differences (Rosenthal, 1991, 1994).

In addition, we compared the differential impact of well-being and ill-being on health outcomes. Effect sizes were similar (though, as expected, in opposite directions), with higher levels of well-being more likely to result in enhanced functioning and higher levels of ill-being more likely to result in compromised functioning. The similar magnitude of these effects sizes was consistent across both experimental and longitudinal study designs. Further, the magnitude of the well-being and ill-being relations was relatively consistent across all three general health outcomes – that is, short-term outcomes, long-term outcomes, and disease/symptom control. Thus, these results demonstrate that the effect of SWB on health is not solely due to ill-being having a detrimental impact on health, but also to well-being having a salutary impact on health. Future research should seek to extend these findings on the differential impact of ill-being and well-being, with a focus on the fundamental underlying mechanisms involved.

Influential moderators

An important benefit of a meta-analytic study is that its method of synthesizing primary results allows for statistical testing of moderators to determine the factors that affect the relations under investigation (Rosenthal & DiMatteo, 2001). In the present analysis, we examined the impact of several factors, including study design, health outcome, the effect of state *vs.* trait well-being, and sample characteristics.

Study design. First, our analyses clearly showed that study design was an important moderator, with experimental studies showing the strongest effects, as expected. This result corroborates the pattern of findings documented by Lyubomirsky et al. (2005), who reported the highest r s for the effects of experimentally induced positive affect on a variety of outcomes. To confirm that this finding was not simply a function of including similar studies as Lyubomirsky et al., we conducted a follow-up analysis in which we considered only the studies not originally included in that review. This follow-up analysis, which examined 80 new experiments, confirmed the larger effect sizes for experimental studies. There may be several reasons that experimental manipulations of affect produce the

biggest effects on health, the most likely being the control that experimental studies have over extraneous variables. In longitudinal and ambulatory studies, other factors, such as psychosocial attributes and measurement differences, are likely to play a relatively greater role (Nesselroade, 1988).

Ambulatory studies offer an interesting cross between short-term processes (observed in experimental manipulations) that are embedded within more long-term periods (Little, Bovaird, & Slegers, 2006). Across the different well-being predictors and outcomes, studies that used ambulatory procedures had significantly lower effect sizes (most of which were not significantly different from zero) than those that used either longitudinal or experimental procedures. We considered the possibility that the ambulatory studies all focused on a single health outcome by examining the effect sizes of the ambulatory studies for the different health outcomes. Ambulatory studies examined eight different health outcomes and only one health outcome (respiratory diseases/conditions) was significantly associated with well-being ($r_{random} = -0.166$). All other health outcomes were non-significant and near zero. Further, the effect sizes within each health outcome typically varied when experimental and ambulatory studies were directly compared. For example, experimental studies that assessed immune functioning reported a large positive association with well-being ($r_{random} = 0.371$), whereas ambulatory studies reported the same association to be near zero ($r_{random} = 0.021$).

Thus, it may be that the transient emotions in day-to-day life that are typically measured in ambulatory studies are simply more readily influenced by other variables (such as the weather, time of day, or daily hassles) that attenuate associations between such emotions and health. Alternatively, to date many fewer studies have used ambulatory methodology, and thus researchers are still establishing the best well-being measures to use and appropriate statistical techniques to analyze such data (Little et al., 2006; Mroczek et al., 2006). It will be important in the future to determine how health outcomes should be conceptualized and measured, how much our measures of health outcomes can be extended across laboratory and field settings and across short- and long-term measurement occasions, and the appropriate statistical techniques that should be used in considering such relations (Mroczek et al., 2006).

Health outcomes. One of the main goals of this meta-analysis was to examine the impact of well-being on specific health outcomes. We expected differential effects depending on whether health was defined in terms of short-term states or as long-term processes, predicting that well-being would most strongly impact short-term outcomes. The data support this hypothesis; however, the stronger short-term well-being–health relations were due to strong associations between well-being and immune functioning and pain tolerance. Short-term effects may be more directly observable, as fewer intervening variables impact the effects that researchers observe; over time, multiple, complex factors potentially moderate long-term health outcomes (Friedman, 2007; Hall et al., 1994). Further, it has been suggested that, at the molecular level, well-being may improve health more

directly both by enhancing immune system response and buffering the system from negative effects of stress (Pressman & Cohen, 2005; Smith, 2006).

This hypothesis can be examined by comparing the average effect sizes observed between well-being and increased immune functioning and pain tolerance with the average effect sizes between well-being and endocrine system response, cardiovascular reactivity, and physiological response. The average effect size for the 69 samples that measured immune functioning and pain tolerance was dramatically higher ($r_{random} = 0.316$) than the 80 studies that measured endocrine system response, cardiovascular reactivity, and physiological response ($r_{random} = -0.009$). These results suggest that rather than buffering from cardiovascular and endocrine system response, well-being may be more likely to lead to a rapid recovery to baseline after a stressor is experienced. As an increase in cardiovascular and endocrine activity is a normal response to stress (e.g., Kemeny, 2007), well-being may counter chronic system activation rather than interrupt normal functioning. This finding corroborates the findings of Fredrickson and her colleagues (see Fredrickson & Levenson, 1998; Fredrickson, Mancuso, Branigan, & Tugade, 2000), who have found that people's cardiovascular activation (after a stressful situation) returns more quickly to their baseline levels after watching positive emotion-inducing films. Further, it appears that well-being not only aids in deregulation of the ANS, but also increases immune response; thus, well-being may affect multiple biological processes.

In addition, some of the most commonly assessed physiological markers had the strongest associations with well-being. For example, the relation between transient positive emotions and sIgA antibody production was the single strongest well-being–health effect size in the meta-analysis. This strong relation may be due to the ease at which sIgA antibody production is measured (through saliva), but future work should aim to determine why positive emotions exert such a strong influence on this immune response. Similarly, positive emotions produce a significant drop in cortisol, but a non-significant drop in all other stress hormones. The stronger effect may be due to cortisol being the stress hormone that is most susceptible to emotional triggers (Dickerson & Kemeny, 2004), whereas other stress hormones, such as epinephrine and norepinephrine, are activated by other types of triggers (such as physical forms of stress). Further, the results of the 32 studies that measured blood pressure revealed a marginally significant, positive association with well-being. A secondary analysis of these data demonstrated that whereas blood pressure increased as a result of increased positive emotions ($r_{random} = 0.153$), blood pressure increased more in the presence of negative emotions. Thus, while positive emotions do result in increases in blood pressure, these increases are smaller than the increases observed for negative emotions.

Operationalizations of well-being. Furthermore, we examined which health outcomes were most strongly associated with state and trait measures of well-being. We expected that short-term health outcomes would be more strongly associated with state manipulations of well-being, whereas long-term health outcomes would

1100 be more strongly associated with trait measures of well-being. With the exception
of the relations between well-being and endocrine response and cardiovascular/
physiological reactivity, this prediction was supported. That transient emotions
have stronger relations with short-term outcomes (especially immune functioning
and pain tolerance) and trait levels of well-being have stronger relations with long-
term outcomes is informative to future investigations. Researchers interested in
altering short-term health outcomes (such as infections or immune system
response) may need to focus on increasing transient emotions, whereas
1105 researchers interested in modifying long-term health outcomes (such as cardio-
vascular outcomes or survival) may need to focus on improving more general
cognitive assessments of well-being.

Sample specific moderators. In addition, we examined specific sample character-
istics, including initial health status, age, and gender composition of the samples.
We expected well-being to have a greater impact where dysregulation is typically
1110 more evident, such as in unhealthy individuals and older individuals (Solomon &
Benton, 1994). Results offered mixed support for these hypotheses. For initial
health status, well-being had a greater impact for both short- and long-term
outcomes in healthy samples; however, well-being more strongly impacted
unhealthy samples in controlling disease and increasing survival. This suggests
1115 that for healthy samples, well-being may enhance functioning at both molecular
and molar levels, whereas for unhealthy samples, well-being may buffer from
subsequent decline. This is important because it suggests that promoting well-
being may indeed help bring about better physical functioning (Rowe, 1988),
especially for healthy individuals, and may improve symptom control for
1120 unhealthy individuals. Considering that happiness interventions have demon-
strated that individuals can increase their well-being by triggering “upward
spirals” through the practice of specific daily behaviors (cf. Lyubomirsky,
Sheldon, & Schkade, 2005; Seligman, Steen, Park, & Peterson, 2005), the
most successful positive psychological interventions may be those that ultimately
1125 increase the health of physically well individuals, and decrease the disease
progression of already physically ill individuals.

Contrary to our predictions, the effects of well-being were fairly constant across
age and gender. However, it is interesting to note where these characteristics did
make a difference. Age moderated the link between well-being and cardiovascular
and physiological reactivity. As individuals age, the risk of cardiovascular-related
1130 incidents significantly increases (Siegler, Bosworth, & Elias, 2003; Siegler, Poon,
Madden, & Dilworth-Anderon, 2004). Also, chronic stress on the cardiovascular
system over time may increase strain on the heart and lead to heart-related
problems (Cacioppo & Berntson, 2007). Although the exact mechanisms are
unclear, that well-being did decrease reactivity implies that well-being may indeed
1135 act as a buffer from strain on the system (McEwen, 1998) for individuals as they
age. Further, males showed a stronger effect in long-term functioning outcomes.
Through both direct and indirect pathways (Cacioppo & Berntson, 2007;
Pressman & Cohen, 2005), well-being may compensate for other vulnerabilities

1140 that have led to greater mortality risk for males. Thus, well-being may play a more
important role for males. Future research should include these characteristics, as
age and gender differences may help us better understand the mechanisms linking
1155 well-being and health outcomes.

Limitations

1145 Any research synthesis is only as good as the current research available to be
meta-analytically combined. Thus, this meta-analysis is limited by the specific
variables that were not manipulated, measured, or reported. First, future
researchers may want to examine those specific health outcomes that have been
1150 studied with relative infrequency. For example, only three relatively small
experimental studies have investigated the impact of well-being on decreased
allergy symptoms, and only eight studies assessed the effect of well-being on the
rate of disease progression.

Second, most of the studies included in our meta-analysis focused on healthy
populations – either students or healthy community members. This was
1155 especially true for the health outcomes related to immune functioning, endocrine
functioning, and cardiovascular and physiological reactivity (all of which are likely
important outcomes for any unhealthy sample). Although evidence is mounting
that higher level of quality of life in is predictive of survival for unhealthy samples
(e.g., cancer patients; see Gotay, 2006), the quality of life measures that have
1160 been used with these unhealthy samples have typically focused on physical
symptoms and health problems rather than on emotional responses or global
judgments of life satisfaction. Thus, future research needs to focus on examining
the impact of hedonic well-being on health for unhealthy participants with a
variety of conditions.

1165 Third, adolescent samples were scarce in the reported research, and, as a result,
any current conclusions about the effects of well-being on health for children and
adolescents would be tenuous at this time. Finally, many studies did not report
information regarding the ethnicity and marital status of their participants.
Therefore, these analyses could not be conducted. Given that these variables may
1170 moderate many of the relations between well-being and health, we encourage
researchers to report these descriptive statistics with greater frequency.

Conclusions

The purpose of this meta-analysis was to examine the effect of well-being on
health outcomes. Much of the previous literature has focused either on the strong
1175 relation between ill-being and health or on how health influences well-being. Our
findings compliment these other studies. Not only can health impact well-being,
as has been established in many other investigations, but well-being can also
impact health. Furthermore, extending earlier research, our analyses highlight the
complex interrelations between well-being and health. Notably, our findings

1180 point to potential biological pathways, such that well-being can directly bolster
immune functioning and buffer the impact of stress.

1185 That well-being can affect short-term and long-term health outcomes and
buffer decline in disease is informative for potential medical and psychological
interventions. Health has been a primary concern throughout history (Ryff &
Singer, 1998), and our findings suggest that a prime area for health promotion
involves boosting happiness and increasing the frequency of positive emotions.
Indeed, health may be only one of many life domains – albeit a critical one – that
is impacted when people actively enhance their own well-being (Lyubomirsky
et al., 2005). Furthermore, from a public health standpoint, mortality and
1190 morbidity are important (Fries, 1990; Kaplan, 2003). As morbidity increases,
health care utilization increases, which in turn escalates health care costs. This
escalation is a problem that pervades the U.S. health care system (Friedman,
1991; Kaplan, 2003; Ryff & Singer, 1998). Thus, to address the question, “What
are the benefits of well-being?” we conclude that the benefits extend from
1195 individuals’ health to a society’s health care costs. Accordingly, the problem of
how to increase and sustain happiness should continue to be pursued by positive
psychologists and health psychologists alike.

Acknowledgements

1200 We are appreciative of the numerous comments and edits on early drafts of this
paper by Colleen J. Howell, Ph.D. We are also grateful to Danielle O’Brien,
Yazmin Perez, and Katrina Rodzon for assistance with preparing the manuscript.

Notes

1205 ¹ For the studies included in the meta-analysis, experimental investigations typically followed a
similar paradigm. Well-being and physiological variables were measured at baseline, mood or
emotion was manipulated, and the physiological variables were measured one or more times; mood/
emotion was again assessed immediately following a manipulation check, and the physiological
variables were again assessed. In some experiments, subjects acted as their own control,
experiencing each mood condition; their reactivity in each condition was compared across
1210 conditions and to their baseline level, using repeated measures analysis of variance or similar
methods (e.g., Brosschot & Thaler, 2003; Clark, Iverson, & Goodwin, 2001; Codispoti et al.,
2003).

1215 Other studies randomly assigned participants to a single mood/emotion condition and
compared between subjects, either controlling for baseline levels or using change scores (e.g.,
Gendolla & Krüsken, 2001a, b). Some of the more recent studies have incorporated multi-level
modeling methods to analyze within and between person changes (e.g., Polk, Cohen, Doyle,
Skoner, & Kirschbaum, 2005). We do note that there is a lot of variation by study, depending on the
outcome of interest, the size of the sample, and the methods used. For example, in one study,
1220 pictures were used to induce positive, negative, or neutral moods (Codispoti et al., 2003). Ten
participants experienced each condition, 1 week apart, in counterbalanced order. Blood was drawn
at baseline, 30 min after baseline, and after the manipulation, and one-way repeated multivariate
analysis of variance was used to analyze the effect of picture valance on several neuroendocrine
markers. In another study, 54 students were randomly assigned to a negative or positive mood, and
to an easy or difficult task (Gendolla & Krüsken, 2001a). Heart rate, blood pressure, and skin
conductance were continually monitored. Change scores between the average baseline function,

during manipulation, and post-manipulation were used to assess the effect of mood on physiological response.

² The most straightforward way to interpret the meta-regression statistics from Table VII is to write out the regression equations from these tables. The basic equation would be as follows:

$$* = \beta_0 + \beta_1$$

where * is the predicted effect size; β_0 is the intercept (when $\beta_1 = 0$); and β_1 is the slope (the change in the predicted effect size with a unit change in the predictor). For example, if we consider the first significant slope from Table VII (percentage of male respondents), we can use the meta-regression coefficients to predict the effect size between well-being and improved immune functioning for samples differing in gender composition. In this case, the predictor (gender composition) runs from 0.00 (a completely female sample) to 1.00 (a completely male sample). Thus, the regression equation for this example (see Table VII) would be

$$* \text{ (predicted effect size)} = 0.392 + -0.267(X_1)$$

where X_1 is the proportion of the sample that is male. For example, if the proportion is 0.00 (completely female sample), the predicted effect size between well-being and improved immune functioning is 0.392. If the proportion is 0.50 (half female/half male), the predicted effect size is 0.258. If the proportion is 1.00 (completely male sample), the predicted effect size is 0.125. We observe that as the sample composition becomes more dominated by males, the strength of the well-being-improved health effect size decreases. These data suggest that well-being may be more strongly related to improved immune functioning for females.

References

*References marked with an asterisk indicate studies included in the meta-analysis.

- Aaronson, N. K., Ahmedzai, S., Bergman, B., Bullinger, M., Cull, A., Duez, N. J. et al. (1993). The European Organization for Research and Treatment of Cancer QLQ-30: A quality-of-life instrument for use in international clinical trials in oncology. *Journal of the National Cancer Institute*, 85, 365–376.
- *Affleck, G., Apter, A., Tennen, H., Reisine, S., Barrows, E., Willard, A. et al. (2000). Mood states associated with transitory changes in asthma symptoms and peak expiratory flow. *Psychosomatic Medicine*, 62, 61–68.
- *Alden, A. L., Dale, J. A., & DeGood, D. E. (2001). Interactive effects of the affect quality and directional focus of mental imagery on pain analgesia. *Applied Psychophysiology and Biofeedback*, 26, 117–126.
- Aldwin, C. M., Spiro, A. III, Levenson, M. R., & Cupertino, A. P. (2001). Longitudinal findings from the normative aging study: III. Personality, individual health trajectories, and mortality. *Psychology and Aging*, 16, 450–465.
- *Apter, A. J., Affleck, G., Reisine, S. T., Tennen, H. A., Barrows, E., Wells, M. et al. (1997). Perception of airway obstruction in asthma: Sequential daily analyses of symptoms, peak expiratory flow rate, and mood. *Journal of Allergy and Clinical Immunology*, 99, 605–612.
- *Avia, M. D., & Kanfer, F. H. (1980). Coping with aversive stimulation: The effects of training in a self-management context. *Cognitive Therapy & Research*, 4, 73–81.
- Bachen, E. A., Manuck, S. B., Marsland, A. L., Cohen, S., Malkoff, S. B., Muldoon, M. F. et al. (1992). Lymphocyte subsets and cellular immune responses to a brief experimental stressor. *Psychosomatic Medicine*, 54, 673–679.
- *Bacon, S. L., Watkins, L. L., Babyak, M., Sherwood, A., Hayano, J., Hinderliter, A. L. et al. (2004). Effects of daily stress on autonomic cardiac control in patients with coronary artery disease. *American Journal of Cardiology*, 93, 1292–1294.
- Baltes, P. B., Staudinger, U. M., & Lindenberger, U. (1999). Lifespan psychology: Theory and application to intellectual functioning. *Annual Review of Psychology*, 50, 471–507.

Well-being and health: A meta-analysis 43

- Berg, C. J., & Snyder, C. R. (2006, October). *The effectiveness of a hope intervention in coping with cold pressor pain*. Paper presented at the 2006 International Positive Psychology Summit, Washington, DC.
- *Berk, L. S., Tan, S. A., Fry, W. F., Napier, B. J., Lee, J. W., Hubbard, R. W. et al. (1989). Neuroendocrine and stress hormone changes during mirthful laughter. *American Journal of the Medical Sciences*, 298, 390–396.
- *Berk, L. S., Felten, D. L., Tan, S. A., Bittman, B. B., & Westengard, J. (2001). Modulation of neuroimmune parameters during the eustress of humor-associated mirthful laughter. *Alternative Therapies in Health and Medicine*, 7, 62–76.
- Bilder, G. E. (1975). Studies on immune competence in the rat: Changes with age, sex, and strain. *Journal of Gerontology*, 30, 641–646.
- AQ1** Boehm, J. K., & Lyubomirsky, S. (in press). Enduring happiness. In S. J. Lopez (Ed.), *Handbook of positive psychology*. Oxford: Oxford University Press.
- *Boiten, F. (1996). Autonomic response patterns during voluntary facial action. *Psychophysiology*, 33, 123–131.
- Booth-Kewley, S., & Friedman, H. S. (1987). Psychological predictors of heart disease: A quantitative review. *Psychological Bulletin*, 101, 343–362.
- Borenstein, M., Hedges, L., Higgins, J., & Rothstein, H. (2005). *Comprehensive meta-analysis, Version 2*. Englewood, NJ: Biostat.
- Bowen, R. A. (2001). Control of endocrine activity. Pathophysiology of the endocrine system. Retrieved November 14, 2006, from www.vivo.colostate.edu/hbooks/pathphys/endocrine/basics/control.html
- Bradburn, N. M. (1969). *The structure of psychological well-being*. Oxford: Aldine.
- Bradburn, N., & Caplovitz, D. (1965). *Reports of happiness*. Chicago, IL: Aldine.
- Breslow, L. (1972). A quantitative approach to the World Health Organization definition of health: Physical, mental and social well-being. *International Journal of Epidemiology*, 1, 347–355.
- *Brosschot, J. F., & Thayer, J. F. (2003). Heart rate response is longer after negative emotions than after positive emotions. *International Journal of Psychophysiology*, 50, 181–187.
- *Brown, J. E., Butow, P. N., Culjak, G., Coates, A. S., & Dunn, S. M. (2000). Psychosocial predictors of outcome: Time to relapse and survival in patients with early stage melanoma. *British Journal of Cancer*, 83, 1448–1453.
- *Brown, W. A., Sirota, A. D., Niaura, R., & Engebretson, T. O. (1993). Endocrine correlates of sadness and elation. *Psychosomatic Medicine*, 55, 458–467.
- *Bruehl, S., Carlson, C. R., & McCubbin, J. A. (1993). Two brief interventions for acute pain. *Pain*, 54, 29–36.
- *Buchanan, T. W., al'Absi, M., & Lovallo, W. R. (1999). Cortisol fluctuates with increases and decreases in negative affect. *Psychoneuroendocrinology*, 24, 227–241.
- Cacioppo, J. T., & Tassinary, L. G. (1990). Inferring psychological significance from physiological signals. *American Psychologist*, 45, 16–28.
- Cacioppo, J. T., & Berntson, G. G. (2007). The brain, homeostasis, and health: Balancing demands of the internal and external milieu. In H. S. Friedman & R. C. Silver (Eds), *Foundations of health psychology* (pp. 73–91). New York, NY: Oxford University Press.
- Cacioppo, J. T., Gardner, W. L., & Berntson, G. G. (1999). The affect system has parallel and integrative processing components: Form follows function. *Journal of Personality and Social Psychology*, 76, 839–855.
- Cannon, W. B. (1932). *Wisdom of the body*. New York, NY: Norton.
- *Carson, M. A., Hathaway, A., Tuohey, J. P., & McKay, B. M. (1988). The effect of a relaxation technique on coronary risk factors. *Behavioral Medicine*, 14, 71–77.
- *Carter, L. E., McNeil, D. W., Vowles, K. E., Sorrell, J. T., Turk, C. L., Ries, B. J. et al. (2002). Effects of emotion on pain reports, tolerance and physiology. *Pain Research & Management*, 7, 21–30.
- Carver, C. S. (2007). Stress, coping, and health. In H. S. Friedman & R. C. Silver (Eds), *Foundations of health psychology* (pp. 117–144). New York, NY: Oxford University Press.
- *Cassileth, B. R., Lusk, E. J., Miller, D. S., Brown, L. L., & Miller, C. (1985). Psychosocial correlates of survival in advanced malignant disease? *New England Journal of Medicine*, 312, 1551–1555.

- Cella, D. F., Tulsky, D. S., Gray, G., Sarafian, B., Linn, E., Bonomi, A., et al. (1993). The Functional Assessment of Cancer Therapy scale: Development and validation of the general measure. *Journal of Clinical Oncology*, *11*, 570–579.
- *Christie, I. C., & Friedman, B. H. (2004). Autonomic specificity of discrete emotion and dimensions of affective space: A multivariate approach. *International Journal of Psychophysiology*, *51*, 143–153.
- *Clark, L., Iversen, S. D., & Goodwin, G. M. (2001). The influence of positive and negative mood states on risk taking, verbal fluency, and salivary cortisol. *Journal of Affective Disorders*, *63*, 179–187.
- Clipp, E. C., Pavalko, E. K., & Elder, G. H. Jr. (1992). Trajectories of health: In concept and empirical pattern. *Behavior, Health, and Aging*, *2*, 159–179.
- Coan, J. A., & Allen, J. J. B. (Eds). (2007). *Handbook of emotion elicitation and assessment*. New York, NY: Oxford University Press.
- *Codispoti, M., Gerra, G., Montebanocci, O., Zaimovic, A., Raggi, M. A., & Baldaro, B. (2003). Emotional perception and neuroendocrine changes. *Psychophysiology*, *40*, 863–868.
- *Cogan, R., Cogan, D., Waltz, W., & McCue, M. (1987). Effects of laughter and relaxation on discomfort thresholds. *Journal of Behavioral Medicine*, *10*, 139–144.
- Cohen, S. (1994). Psychosocial influences on immunity and infectious disease in humans. In R. Glaser & J. Kiecolt-Glaser (Eds), *Handbook of human stress and immunity* (pp. 301–319). San Diego, CA: Academic Press.
- Cohen, S., & Williamson, G. M. (1991). Stress and infectious disease in humans. *Psychological Bulletin*, *109*, 5–24.
- *Cohen, S., Doyle, W. J., Turner, R. B., Alper, C. M., & Skoner, D. P. (2003). Emotional style and susceptibility to the common cold. *Psychosomatic Medicine*, *65*, 652–657.
- Cooper, H. (1998). *Synthesizing research: A guide for literature reviews* (3rd ed.). Beverly Hills, CA: Sage.
- Curran, S. L., Andrykowski, M. A., & Studts, J. L. (1995). Short form of the Profile of Mood States (POMS-SF): Psychometric information. *Psychological Assessment*, *7*, 80–83.
- *Danner, D. D., Snowdon, D. A., & Friesen, W. V. (2001). Positive emotions in early life and longevity: Findings from the nun study. *Journal of Personality and Social Psychology*, *80*, 804–813.
- *Davidson, R. J., Kabat-Zinn, J., Schumacher, J., Rosenkranz, M., Muller, D., Santorelli, S. F., et al. (2003). Alternations in brain and immune function produced by mindfulness meditation. *Psychosomatic Medicine*, *65*, 564–570.
- Dayyani, F., Joeinig, A., Ziegler-Heitbrock, L., Schmidmaier, R., Straka, C., Emmerich, B., et al. (2004). Autologous stem-cell transplantation restores the functional properties of CD14+ CD16+ monocytes in patients with myeloma and lymphoma. *Journal of Leukocyte Biology*, *75*, 207–213.
- *Deeg, D. J. H., & Van Zonneveld, R. J. (1989). Does happiness lengthen life? The prediction of longevity in the elderly. In R. Veenhoven (Ed.), *How harmful is happiness? Consequences of enjoying life or not* (pp. 29–34). Rotterdam: Universitaire Pers Rotterdam.
- *Derogatis, L. R., Abeloff, M. D., & Melisaratos, N. (1979). Psychological coping mechanisms and survival time in metastatic breast cancer. *Journal of the American Medical Association*, *242*, 1504–1508.
- AQ2** *Devins, G. M., Mann, J., Mandin, H. P., & C., L. (1990). Psychosocial predictors of survival in end-stage renal disease. *Journal of Nervous and Mental Disease*, *178*, 127–133.
- Dickerson, S. S., & Kemeny, M. E. (2004). Acute stressors and cortisol responses: A theoretical integration and synthesis of laboratory research. *Psychological Bulletin*, *130*, 355–391.
- Diener, E. (1994). Assessing subjective well-being: Progress and opportunities. *Social Indicators Research*, *31*, 103–157.
- Diener, E. (2000). Subjective well-being: The science of happiness and a proposal for a national index. *American Psychologist*, *55*, 34–43.
- Diener, E., & Emmons, R. A. (1984). The independence of positive and negative affect. *Journal of Personality and Social Psychology*, *47*, 1105–1117.
- Diener, E., Emmons, R. A., Larsen, R. J., & Griffin, S. (1985). The satisfaction with life scale. *Journal of Personality Assessment*, *49*, 71–75.

Well-being and health: A meta-analysis 45

- 1385 Diener, E., Sandvik, E., & Pavot, W. (1991). Happiness is the frequency, not the intensity, of positive versus negative affect. In F. Strack, M. Argyle & N. Schwarz (Eds), *Subjective well-being: An interdisciplinary perspective* (pp. 119–139). New York, NY: Pergamon.
- Diener, E., Smith, H. L., & Fujita, F. (1995). The personality structure of affect. *Journal of Personality and Social Psychology*, 69, 130–141.
- 1390 Diener, E., Suh, E. M., Lucas, R. E., & Smith, H. L. (1999). Subjective well-being: Three decades of progress. *Psychological Bulletin*, 125, 276–302.
- *Dillon, K. M., Minchoff, B., & Baker, K. H. (1985). Positive emotional states and enhancement of the immune system. *International Journal of Psychiatry in Medicine*, 15, 13–18.
- Dreher, H. (1995). The immune system: A primer. In H. Dreher (Ed.), *The immune power personality*. Retrieved November 13, 2006, from www.thebody.com/dreher/primer.html
- 1395 *Ekman, P., Levenson, R. W., & Friesen, W. V. (1983). Autonomic nervous system activity distinguishes among emotions. *Science*, 221, 1208–1210.
- AQ3** *Evans, P., Bristow, M., Hucklebridge, F., Clow, A. et al. (1993). The relationship between secretory immunity, mood and life-events. *British Journal of Clinical Psychology*, 32, 227–236.
- 1400 Fisher, B. J. (1995). Successful aging, life satisfaction, and generativity in later life. *International Journal of Aging and Human Development*, 41, 239–250.
- *Florin, I., Freudenberg, G., & Hollaender, J. (1985). Facial expressions of emotion and physiologic reactions in children with bronchial asthma. *Psychosomatic Medicine*, 47, 382–393.
- *Foster, P. S., Webster, D. G., & Williamson, J. (2003). The psychophysiological differentiation of actual, imagined, and recollected mirth. *Cognition and Personality*, 22, 163–180.
- 1405 *Frazier, T. W., Strauss, M. E., & Steinhauer, S. R. (2004). Respiratory sinus arrhythmia as an index of emotional response in young adults. *Psychophysiology*, 41, 75–83.
- *Fredrickson, B. L., & Levenson, R. W. (1998). Positive emotions speed recovery from the cardiovascular sequelae of negative emotions. *Cognition and Emotion*, 12, 191–220.
- 1410 *Fredrickson, B. L., Mancuso, R. A., Branigan, C., & Tugade, M. M. (2000). The undoing effect of positive emotions. *Motivation and Emotion*, 24, 237–258.
- Friedman, H. S. (1991). *The self-healing personality*. New York, NY: Henry Holt.
- Friedman, H. S. (2007). Personality, disease, and self-healing. In H. S. Friedman & R. C. Silver (Eds), *Foundations of Health Psychology* (pp. 172–199). New York, NY: Oxford University Press.
- 1415 Friedman, H. S., & Booth-Kewley, S. (1987). The “disease-prone personality”: A meta-analytic view of the construct. *American Psychologist*, 42, 539–555.
- AQ4** *Friedman, H. S., Tucker, J. S., Tomlinson-Keasey, C., Schwartz, J. E., Wingard, & Criqui, M. H. (1993). Does childhood personality predict longevity? *Journal of Personality and Social Psychology*, 65, 176–185.
- 1420 Fries, J. F. (1990). Medical perspectives upon successful aging. In P. B. Baltes & M. M. Baltes (Eds), *Successful aging: Perspectives from the behavioral sciences* (pp. 35–49). New York, NY: Cambridge University Press.
- *Futterman, A. D., Kemeny, M. E., Shapiro, D., Polonsky, W., & Fahey, J. L. (1992). Immunological variability associated with experimentally-induced positive and negative affective states. *Psychological Medicine*, 22, 231–238.
- 1425 *Futterman, A. D., Kemeny, M. E., Shapiro, D., & Fahey, J. L. (1994). Immunological and physiological changes associated with induced positive and negative mood. *Psychosomatic Medicine*, 56, 499–511.
- *Gellman, M., Spitzer, S., Ironson, G., Llabre, M., Saab, P., DeCarlo Pasin, R. et al. (1990). Posture, place, and mood effects on ambulatory blood pressure. *Psychophysiology*, 27, 544–551.
- *Gendolla, G. H., & Krusken, J. (2001a). The joint impact of mood state and task difficulty on cardiovascular and electrodermal reactivity in active coping. *Psychophysiology*, 38, 548–556.
- 1435 *Gendolla, G. H., & Krusken, J. (2001b). Mood state and cardiovascular response in active coping with an affect-regulative challenge. *International Journal of Psychophysiology*, 41, 169–180.
- *Gendolla, G. H., & Krusken, J. (2002). Mood state, task demand, and effort-related cardiovascular response. *Cognition & Emotion*, 16, 577–603.
- *Gendolla, G. H. E., Abele, A. E., & Krusken, J. (2001). The informational impact of mood on effort mobilization: A study of cardiovascular and electrodermal responses. *Emotion*, 1, 12–24.
- 1440 *Giltay, E. J., Geleijnse, J. M., Zitman, F. G., Hoekstra, T., & Schouten, E. G. (2004). Dispositional optimism and all-cause and cardiovascular mortality in a prospective cohort of elderly Dutch men and women. *Archives of General Psychiatry*, 61, 1126–1135.

- Glaser, R., & Kiecolt-Glaser, J. (Eds). (1994). *Handbook of human stress and immunity*. San Diego, CA: Academic Press.
- Gochman, D. S. (1997). Health behavior research: Definitions and diversity. In D. S. Gochman (Ed.), *Handbook of healthy behavior research I: Personal and social determinants* (pp. 3–20). New York, NY: Plenum Press.
- *Gomez, P., Zimmermann, P., Guttormsen-Schär, S., & Danuser, B. (2005). Respiratory responses associated with affective processing of film stimuli. *Biological Psychology*, 68, 223–235.
- Gotay, C. (2006, October). *The prognostic value of QOL ratings in cancer patients: A systematic review*. Paper presented at the 13th Annual Conference of the International Society for Quality of Life Research, Lisbon, Portugal.
- Grzywacz, J. G., & Keyes, C. L. M. (2004). Toward health promotion: Physical and social behaviors in complete health. *American Journal of Health Behaviors*, 28, 99–111.
- *Gullette, E. C., Blumenthal, J. A., Babyak, M., Jiang, W., Waugh, R. A., Frid, D. J., et al. (1997). Effects of mental stress on myocardial ischemia during daily life. *Journal of the American Medical Association*, 277, 1521–1526.
- Guralnik, J. M., & Kaplan, G. A. (1989). Predictors of healthy aging: Prospective evidence from the Alameda County Study. *American Journal of Public Health*, 79, 703–708.
- Hall, N. R. S., Anderson, J. A., & O’Grady, M. P. (1994). Stress and immunity in humans: Modifying variables. In R. Glaser & J. Kiecolt-Glaser (Eds), *Handbook of human stress and immunity* (pp. 183–215). San Diego, CA: Academic Press.
- Hampson, S. E., Goldberg, L. R., Vogt, T. M., & Dubanoski, J. P. (2006). Forty years on: Teachers’ assessments of children’s personality traits predict self-reported health behaviors and outcomes at midlife. *Health Psychology*, 25, 57–64.
- Harker, L., & Keltner, D. (2001). Expressions of positive emotions in women’s college yearbook pictures and their relationship to personality and life outcomes across adulthood. *Journal of Personality and Social Psychology*, 80, 112–124.
- *Harrison, L. K., Carroll, D., Burns, V. E., Corkill, A. R., Harrison, C. M., Ring, C., et al. (2000). Cardiovascular and secretory immunoglobulin A reactions to humorous, exciting, and didactic film presentations. *Biological Psychology*, 52, 113–126.
- Hedges, L. V. (1994). Fixed effects models. In H. Cooper & H. Hedges (Eds), *The handbook of research synthesis* (pp. 285–299). New York, NY: Russell Sage Foundation.
- Herbert, T. B., & Cohen, S. (1993). Stress and immunity in humans: A meta-analytic review. *Psychosomatic Medicine*, 55, 364–379.
- *Hertel, J. B., & Hekmat, H. M. (1994). Coping with cold-pressor pain: Effects of mood and covert imaginal modeling. *Psychological Record*, 44, 207–220.
- *Hess, U., Kappas, A., McHugo, G. J., Lanzetta, J. T., & Kleck, R. E. (1992). The facilitative effect of facial expression on the self-generation of emotion. *International Journal of Psychophysiology*, 12, 251–265.
- *Horan, J. J., & Dellinger, J. K. (1974). “In vivo” emotive imagery: A preliminary test. *Perceptual & Motor Skills*, 39, 359–362.
- *Houghton, L. A., Calvert, E. L., Jackson, N. A., Cooper, P., & Whorwell, P. J. (2002). Visceral sensation and emotion: A study using hypnosis. *Gut*, 51, 701–704.
- *Hubert, W., & de Jong-Meyer, R. (1990). Psychophysiological response patterns to positive and negative film stimuli. *Biological Psychology*, 31, 73–93.
- *Hubert, W., & de Jong-Meyer, R. (1991). Autonomic, neuroendocrine, and subjective responses to emotion-inducing film stimuli. *International Journal of Psychophysiology*, 11, 131–140.
- *Hubert, W., Moller, M., & de Jong-Meyer, R. (1993). Film-induced amusement changes in saliva cortisol levels. *Psychoneuroendocrinology*, 18, 265–272.
- *Hucklebridge, F., Lambert, S., Clow, A., Warburton, D. M., Evans, P. D., & Sherwood, N. (2000). Modulation of secretory immunoglobulin A in saliva; response to manipulation of mood. *Biological Psychology*, 53, 25–35.
- *Hudak, D. A., Dale, J. A., Hudak, M. A., & DeGood, D. E. (1991). Effects of humorous stimuli and sense of humor on discomfort. *Psychological Reports*, 69, 779–786.
- *Hyland, M. E. (1990). The mood-peak flow relationship in adult asthmatics: A pilot study of individual differences and direction of causality. *British Journal of Medical Psychology*, 63, 379–384.
- Idler, E. L., & Kasl, S. (1991). Health perceptions and survival: Do global evaluations of health status really predict mortality? *Journals of Gerontology*, 46, S55–S65.

Well-being and health: A meta-analysis 47

- *Jacob, R. G., Thayer, J. F., Manuck, S. B., Muldoon, M. F., Tamres, L. K., Williams, D. M. et al. (1999). Ambulatory blood pressure responses and the circumplex model of mood: A 4-day study. *Psychosomatic Medicine*, 61, 319–333.
- Kaplan, R. M. (1994). The Ziggy theorem: Toward an outcomes-focused health psychology. *Health Psychology*, 13, 451–460.
- Kaplan, R. M. (2003). The significance of quality of life in health care. *Quality of Life Research*, 12, 3–16.
- *Kawamoto, R., & Doi, T. (2002). Self-reported functional ability predicts three-year mobility and mortality in community-dwelling older persons. *Geriatrics and Gerontology International*, 2, 68–74.
- Keller, S. E., Shiflett, S. C., Schleifer, S. J., & Bartlett, J. A. (1994). Stress, immunity, and health. In R. Glaser & J. Kiecolt-Glaser (Eds), *Handbook of human stress and immunity* (pp. 217–244). San Diego, CA: Academic Press.
- Kemeny, M. E. (2007). Psychoneuroimmunology. In H. S. Friedman & R. C. Silver (Eds), *Foundations of health psychology* (pp. 92–116). New York, NY: Oxford University Press.
- Kiecolt-Glaser, J. K., Malarkey, W. B., Cacioppo, J. T., & Glaser, R. (1994). Stressful personal relationships: Immune and endocrine function. In R. Glaser & J. Kiecolt-Glaser (Eds), *Handbook of human stress and immunity* (pp. 321–339). San Diego, CA: Academic Press.
- Kim, B. (1998). Socioeconomic status and perception of the quality of life in Korea. *Development and Society*, 27, 1–15.
- *Kimata, H. (2004). Effect of viewing a humorous vs. nonhumorous film on bronchial responsiveness in patients with bronchial asthma. *Physiology & Behavior*, 81, 681–684.
- *Kivimäki, M., Vahtera, J., Elovainio, M., Helenius, H., Singh-Manoux, A., & Pentti, J. (2005). Optimism and pessimism as predictors of change in health after death or onset of severe illness in family. *Health Psychology*, 24, 413–421.
- *Knapp, P. H., Levy, E. M., Giorgi, R. G., Black, P. H., Fox, B. H., & Heeren, T. C. (1992). Short-term immunological effects of induced emotion. *Psychosomatic Medicine*, 54, 133–148.
- Koehler, K. R. (1996). The human cardiovascular system. Retrieved November 14, 2006, from www.rwc.uc.edu/koehler/biophys/3a.html
- *Koivumaa-Honkanen, H., Honkanen, R., Viinamäki, H., Heikkilä, K., Kaprio, J., & Koskenvuo, M. (2000). Self-reported life satisfaction and 20-year mortality in healthy Finnish adults. *American Journal of Epidemiology*, 152, 983–991.
- Kozma, A., & Stones, M. J. (1980). The measurement of happiness: Development of the Memorial University of Newfoundland Scale of Happiness (MUNSH). *Journal of Gerontology*, 35, 906–912.
- Krantz, D. S., & McCeney, M. K. (2002). Effects of psychological and social factors on organic disease: A critical assessment of research on coronary heart disease. *Annual Review of Psychology*, 53, 341–369.
- *Krause, J. S., Sternberg, M., Lottes, S., & Maides, J. (1997). Mortality after spinal cord injury: An 11-year prospective study. *Archives of Physical Medicine and Rehabilitation*, 78, 815–821.
- Kubzansky, L. D., & Kawachi, I. (2000). Going to the heart of the matter: Do negative emotions cause coronary heart disease? *Journal of Psychosomatic Research*, 48, 323–337.
- *Kubzansky, L. D., Sparrow, D., Vokonas, P., & Kawachi, I. (2001). Is the glass half empty or half full? A prospective study of optimism and coronary heart disease in the normative aging study. *Psychosomatic Medicine*, 63, 910–916.
- *Kubzansky, L. D., Wright, R. J., Cohen, S., Weiss, S., Rosner, B., & Sparrow, D. (2002). Breathing easy: A prospective study of optimism and pulmonary function in the normative aging study. *Annals of Behavioral Medicine*, 24, 345–353.
- *Kugler, J., & Kalveram, K. T. (1987). Is salivary cortisol related to mood states and psychosomatic symptoms? In H. Weiner, I. Florin, R. Murison & D. Hellhammer (Eds), *Frontiers of stress research* (pp. 388–391). Kirkland, WA: Hans Huber Publishers.
- *Laidlaw, T. M., Booth, R. J., & Large, R. G. (1994). The variability of type I hypersensitivity reactions: The importance of mood. *Journal of Psychosomatic Research*, 38, 51–61.
- *Laidlaw, T. M., Booth, R. J., & Large, R. G. (1996). Reduction in skin reactions to histamine after a hypnotic procedure. *Psychosomatic Medicine*, 58, 242–248.
- *Lambert, R. B., & Lambert, N. K. (1995). The effects of humor on secretory immunoglobulin A levels in school-aged children. *Pediatric Nursing*, 21, 16–19.

- Laudenslager, M. L., & Fleshner, M. (1994). Stress and immunity: Of mice, monkeys, models, and mechanisms. In R. Glaser & J. Kiecolt-Glaser (Eds), *Handbook of human stress and immunity* (pp. 161–181). San Diego, CA: Academic Press.
- *Lefcourt, H. M., Davidson-Katz, K., & Kueneman, K. (1990). Humor and immune-system functioning. *Humor: International Journal of Humor Research*, 3, 305–321.
- Lerner, J. S., Gonzalez, R. M., Dahl, R. E., Hariri, A. R., & Taylor, S. E. (2005). Facial expressions of emotion reveal neuroendocrine and cardiovascular stress responses. *Biological Psychiatry*, 58, 743–750.
- *Levenson, R. W., Ekman, P., & Friesen, W. V. (1990). Voluntary facial action generates emotion-specific autonomic nervous system activity. *Psychophysiology*, 27, 363–384.
- *Levy, B. R., Slade, M. D., Kunkel, S. R., & Kasl, S. V. (2002). Longevity increased by positive self perceptions of aging. *Journal of Personality and Social Psychology*, 83, 261–270.
- *Levy, S. M., Lee, J., Bagley, C., & Lipman, M. (1988). Survival hazard analysis in first recurrent breast cancer patients: Seven-year follow-up. *Psychosomatic Medicine*, 50, 520–528.
- *Liangas, G., Morton, J. R., & Henry, R. L. (2003). Mirth-triggered asthma: Is laughter really the best medicine? *Pediatric Pulmonology*, 36, 107–112.
- Linnemeyer, P. A. (1993). The immune system: An overview. *The Body: The Complete HIV/AIDS Resource*. Retrieved November 13, 2006, from www.thebody.com/step/immune.html
- Little, T. D., Bovaird, J. A., & Slegers, D. W. (2006). Methods for the analysis of change. In D. K. Mroczek & T. D. Little (Eds), *Handbook of personality development* (pp. 181–211). Mahwah, NJ: Lawrence Erlbaum Associates.
- Lucas, R. E., Diener, E., & Suh, E. M. (1996). Discriminant validity of well-being measures. *Journal of Personality and Social Psychology*, 71, 616–628.
- Lyubomirsky, S., & Lepper, H. S. (1999). A measure of subjective happiness: Preliminary reliability and construct validation. *Social Indicators Research*, 46, 137–155.
- Lyubomirsky, S., King, L. A., & Diener, E. (2005). The benefits of frequent positive affect: Does happiness lead to success? *Psychological Bulletin*, 131, 803–855.
- Lyubomirsky, S., Sheldon, K. M., & Schkade, D. (2005). Pursuing happiness: The architecture of sustainable change. *Review of General Psychology*, 9, 111–131.
- Lyubomirsky, S., Tkach, C., & DiMatteo, M. R. (2006). What are the differences between happiness and self-esteem? *Social Indicators Research*, 78, 363–404.
- *Lutgendorf, S. K., Vitaliano, P. P., Tripp-Reimer, T., Harvey, J. H., & Lubaroff, D. M. (1999). Sense of coherence moderates the relationship between life stress and natural killer cell activity in health older adults. *Psychology and Aging*, 14, 552–563.
- *Maier, H., & Smith, J. (1999). Psychological predictors of mortality in old age. *Journal of Gerontology*, 54, 44–P54.
- Makinodan, T., Hahn, T. J., McDougall, S., Yamaguchi, D. T., Fang, M., & Iida-Klein, A. (1991). Cellular immunosenescence: An overview. *Experimental Gerontology*, 26, 281–288.
- Manuck, S. B., Cohen, S., Rabin, B. S., Muldoon, M. F., & Bachen, E. A. (1991). Individual differences in cellular immune response to stress. *Psychological Science*, 2, 1–5.
- Marks, G. N., & Fleming, N. (1999). Influences and consequences of well-being among Australian young people: 1980–1995. *Social Indicators Research*, 46, 301–323.
- Maynahan, J. A., Brenner, G. J., Cocke, R., Karp, J. D., Breneman, S. M., Dopp, J. M. et al. (1994). Stress-induced modulation of immune function in mice. In R. Glaser & J. Kiecolt-Glaser (Eds), *Handbook of human stress and immunity* (pp. 1–22). San Diego, CA: Academic Press.
- *McClelland, D. C., & Cheriff, A. D. (1997). The immunoenhancing effects of humor on secretory IgA and resistance to respiratory infection. *Psychology and Health*, 12, 329–344.
- *McCraty, R., Atkinson, M., Tiller, W. A., Rein, G., & Watkins, A. D. (1995). The effects of emotions on short-term power spectrum analysis of heart rate variability. *American Journal of Cardiology*, 76, 1089–1093.
- *McCraty, R., Atkinson, M., Rein, G., & Watkins, A. D. (1996). Music enhances the effect of positive emotional states on salivary IgA. *Stress Medicine*, 12, 167–175.
- McEwen, B. S. (1998). Protective and damaging effects of stress mediators. *New England Journal of Medicine*, 338, 171–179.
- McEwen, B. S., & Stellar, E. (1993). Stress and the individual. *Archives of Internal Medicine*, 153, 2093–2101.

Well-being and health: A meta-analysis 49

- 1615 *Meagher, M. W., Arnau, R. C., & Rhudy, J. L. (2001). Pain and emotion: Effects of affective picture modulation. *Psychosomatic Medicine*, 63, 79–90.
- *Meininger, J. C., Liehr, P., Chan, W., Smith, G., & Mueller, W. H. (2004). Developmental, gender, and ethnic group differences in moods and ambulatory blood pressure in adolescents. *Annals of Behavioral Medicine*, 28, 10–19.
- 1620 *Milam, J. E., Richardson, J. L., Marks, G., Kemper, C. A., & McCutchan, A. J. (2004). The roles of dispositional optimism and pessimism in HIV disease progression. *Psychology & Health*, 19, 167–181.
- *Miller, B. D., & Wood, B. L. (1997). Influence of specific emotional states on autonomic reactivity and pulmonary function in asthmatic children. *Journal of the American Academy of Child & Adolescent Psychiatry*, 36, 669–677.
- 1625 *Mittwoch-Jaffe, T., Shalit, F., Srendi, B., & Yehuda, S. (1995). Modification of cytokine secretion following mild emotional stimuli. *Neuroreport*, 6, 789–792.
- *Moskowitz, J. T. (2003). Positive affect predicts lower risk of aids mortality. *Psychosomatic Medicine*, 65, 620–626.
- 1630 Mroczek, D. K., & Spiro, A. III (2005). Change in life satisfaction during adulthood: Findings from the Veterans Affairs Normative Aging Study. *Journal of Personality and Social Psychology*, 88, 189–202.
- Mroczek, D. K., Almeida, D. M., Spiro, A. III, & Pafford, C. (2006). Modeling intraindividual stability and change in personality. In D. K. Mroczek & T. D. Little (Eds), *Handbook of personality development* (pp. 163–180). Mahwah, NJ: Lawrence Erlbaum Associates.
- 1635 Myers, D. G., & Diener, E. (1995). Who is happy? *Psychological Science*, 6, 10–19.
- Nesselroade, J. R. (1988). Sampling and generalizability: Adult development and aging issues examined within the general methodological framework of selection. In K. W. Schaie, R. T. Campbell, W. M. Meredith & S. C. Rawlings (Eds), *Methodological issues in aging research* (pp. 108–121). New York, NY: Springer.
- 1640 Nesselroade, J. R. (1991). Interindividual differences in intraindividual change. In L. M. Collins & J. L. Horn (Eds), *Best methods for the analysis of change* (pp. 92–105). Washington, DC: American Psychological Association.
- *Neumann, S. A., & Waldstein, S. R. (2001). Similar patterns of cardiovascular response during emotional activation as a function of affective valence and arousal and gender. *Journal of Psychosomatic Research*, 50, 245–253.
- 1645 *Njus, D. M., Nitschke, W., & Bryant, F. B. (1996). Positive affect, negative affect, and the moderating effect of writing on sIgA antibody levels. *Psychology & Health*, 12, 135–148.
- *O'Connor, B. P., & Vallerand, R. J. (1998). Psychological adjustment variables as predictors of mortality among nursing home residents. *Psychology and Aging*, 13, 368–374.
- 1650 *Ong, A. D., & Allaire, J. C. (2005). Cardiovascular intraindividual variability in later life: The influence of social connectedness and positive emotions. *Psychology and Aging*, 20, 476–485.
- *Ostir, G. V., Markides, K. S., Black, S. A., & Goodwin, J. S. (2000). Emotional well-being predicts subsequent functional independence and survival. *Journal of the American Geriatrics Society*, 48, 473–478.
- 1655 *Ostir, G. V., Markides, K. S., Peek, M. K., & Goodwin, J. S. (2001). The association between emotional well-being and the incidence of stroke in older adults. *Psychosomatic Medicine*, 63, 210–215.
- *Ostir, G. V., Goodwin, J. S., Markides, K. S., Ottenbacher, K. J., Balfour, J., & Guralnik, J. M. (2002). Differential effects of pre morbid physical and emotional health on recovery from acute events. *Journal of the American Geriatrics Society*, 50, 713–718.
- 1660 *Ostir, G. V., Ottenbacher, K. J., & Markides, K. S. (2004). Onset of frailty in older adults and the protective role of positive affect. *Psychology and Aging*, 19, 402–408.
- *Palmore, E. B. (1969). Predicting longevity: A follow-up controlling for age. *Gerontologist*, 9, 247–250.
- 1665 *Parker, M. G., Thorslund, M., & Nordstrom, M. L. (1992). Predictors of mortality for the oldest old. A 4-year follow-up of community-based elderly in Sweden. *Archives of Gerontology and Geriatrics*, 14, 227–237.
- *Perera, S., Sabin, E., Nelson, P., & Lowe, D. (1998). Increases in salivary lysozyme and IgA concentrations and secretory rates independent of salivary flow rates following viewing of a humorous video. *International Journal of Behavioral Medicine*, 5, 118–228.
- 1670

Pickering, T. (1999). How blood pressure is regulated. *Lifeclinic Health Management Systems*. Retrieved online November 14, 2006, from www.lifeclinic.com/focus/blood/article_View.asp?MessageID=248

*Pitkala, K. H., Laakkonen, M. L., Strandberg, T. E., & Tilvis, R. S. (2004). Positive life orientation as a predictor of 10-year outcome in an aged population. *Journal of Clinical Epidemiology*, *57*, 409–414.

*Polk, D. E., Cohen, S., Doyle, W. J., Skoner, D. P., & Kirschbaum, C. (2005). State and trait affect as predictors of salivary cortisol in healthy adults. *Psychoneuroendocrinology*, *30*, 261–272.

*Pollard, T. M., & Schwartz, J. E. (2003). Are changes in blood pressure and total cholesterol related to changes in mood? An 18-month study of men and women. *Health Psychology*, *22*, 47–53.

Pressman, S. D., & Cohen, S. (2005). Does positive affect influence health? *Psychological Bulletin*, *131*, 925–971.

*Prkachin, K. M., Williams-Avery, R. M., Zwaal, C., & Mills, D. E. (1999). Cardiovascular changes during induced emotion: An application of Lang's theory of emotional imagery. *Journal of Psychosomatic Research*, *47*, 255–267.

*Provost, M. A., & Gouin-Décarie, T. (1979). Heart rate reactivity of 9- and 12-months old infants showing specific emotions in natural setting. *International Journal of Behavioral Development*, *2*, 109–120.

Quanjer, P. H., Tammeling, G. J., Cotes, J. E., Pedersen, O. F., Peslin, R., & Yernault, J. C. (1993). Lung volumes and forced ventilatory flows: Official statement of the European Respiratory Society. *The European Respiratory Journal*, *6*, 5–40.

Quanjer, P. H., Lebowitz, M. D., Gregg, I., Miller, M. R., & Pedersen, O. F. (1997). Peak expiratory flow: Conclusions and recommendations of a working party of the European Respiratory Society. *The European Respiratory Journal*, *10*, 2S–8S.

Rabin, B. S. (1999). *Stress, immune function, and health: The connection*. New York, NY: John Wiley-Liss.

Rabin, B. S., Kusnecov, A., Shurin, M., Zhou, D., & Rasnick, S. (1994). Mechanistic aspects of stressor-induced immune alteration. In R. Glaser & J. Kiecolt-Glaser (Eds), *Handbook of human stress and immunity* (pp. 23–51). San Diego, CA: Academic Press.

*Reynolds, D. K., & Nelson, F. L. (1981). Personality, life situation and life expectancy. *Suicide and Life Threatening Behavior*, *11*, 99–110.

*Rhudy, J. L., Williams, A. E., McCabe, K. M., Nguyen, M. A. T. V., & Rambo, P. (2005). Affective modulation of nociception at spinal and supraspinal levels. *Psychophysiology*, *42*, 579–587.

*Richman, L. S., Kubzansky, L., Maselko, J., Kawachi, I., Choo, P., & Bauer, M. (2005). Positive emotion and health: Going beyond the negative. *Health Psychology*, *24*, 422–429.

*Ritz, T., Steptoe, A., DeWilde, S., & Costa, M. (2000). Emotions and stress increase respiratory resistance in asthma. *Psychosomatic Medicine*, *62*, 401–412.

*Ritz, T., Claussen, C., & Dahme, B. (2001). Experimentally induced emotions, facial muscle activity, and respiratory resistance in asthmatic and non-asthmatic individuals. *British Journal of Medical Psychology*, *74*, 167–182.

*Ritz, T., Thöns, M., Fahrenkrug, S., & Dahme, B. (2005). Airways, respiration, and respiratory sinus arrhythmia during picture viewing. *Psychophysiology*, *42*, 568–578.

Roos, N. P., & Havens, B. (1991). Predictors of successful aging: A twelve-year study of Manitoba elderly. *American Journal of Public Health*, *81*, 63–68.

*Rosenbaum, M. (1980). Individual differences in self-control behaviors and tolerance of painful stimulation. *Journal of Abnormal Psychology*, *89*, 581–590.

Rosenow, E. (2005). Oxygen saturation test. Retrieved November 15, 2006, from the MayoClinic.com website: www.mayoclinic.com/health/oxygen-saturation/AN01016

Rosenthal, R. (1991). *Meta-analytic procedures for social research*. Newbury Park, CA: Sage Publications.

Rosenthal, R. (1994). Parametric measures of effect size. In H. Cooper & L. Hedges (Eds), *The handbook of research synthesis* (pp. 231–244). New York, NY: Russell Sage Foundation.

Rosenthal, R., & DiMatteo, M. R. (2001). Meta-analysis: Recent developments in quantitative methods for literature reviews. *Annual Review of Psychology*, *52*, 59–82.

Rosenthal, R., & Rubin, D. B. (2003). *r*-sub(equivalent): A simple effect size indicator. *Psychological Methods*, *8*, 492–496.

Well-being and health: A meta-analysis 51

- 1730 Rowe, J. W. (1988). Aging reconsidered: Strategies to promote health and prevent disease in old age. *Quarterly Journal of Medicine*, 66, 1–4.
- Rowe, J. W., & Kahn, R. L. (1987). Human aging: Usual and successful. *Science*, 237, 143–149.
- Ryff, C. D., & Singer, B. (1998). The contours of positive human health. *Psychological Inquiry*, 9, 1–28.
- 1735 Saleh, M. N., Goldman, S. J., LoBuglio, A. F., Beall, A. C., Sabio, H., McCord, M. C., et al (1995). CD16+ monocytes in patients with cancer: spontaneous elevation and pharmacologic induction by recombinant human macrophage colony-stimulating factor. *Blood*, 85, 2910–2917.
- Sandvik, E., Diener, E., & Seidlitz, L. (1993). Subjective well-being: The convergence and stability of self-report and non-self-report measures. *Journal of Personality*, 61, 317–342.
- 1740 *Santibáñez, H. G., & Bloch, S. (1986). A qualitative analysis of emotional effector patterns and their feedback. *Pavlovian Journal of Biological Sciences*, 21, 108–116.
- Scheier, M. F., & Carver, C. S. (1985). Optimism, coping, and health: Assessment and implications of generalized outcome expectancies. *Health Psychology*, 4, 219–247.
- 1745 *Scheier, M. F., Matthews, K. A., Owens, J. F., Magovern, G. J., Lefebvre, R. C., Abbott, R. A. et al. (1989). Dispositional optimism and recovery from coronary artery bypass surgery: The beneficial effects on physical and psychological well-being. *Journal of Personality and Social Psychology*, 57, 1024–1040.
- Schultz, R., & Heckhausen, J. (1996). A life span model of successful aging. *American Psychologist*, 51, 702–714.
- 1750 *Schwartz, G. E., Weinberger, D. A., & Singer, J. A. (1981). Cardiovascular differentiation of happiness, sadness, anger, and fear following imagery and exercise. *Psychosomatic Medicine*, 43, 343–364.
- *Schwartz, J. E., Warren, K., & Pickering, T. G. (1994). Mood, location and physical position as predictors of ambulatory blood pressure and heart rate: Application of a multi-level random effects model. *Annals of Behavioral Medicine*, 16, 210–220.
- 1755 *Scott, D. S., & Barber, T. X. (1977). Cognitive control of pain: Effects of multiple cognitive strategies. *Psychological Record*, 27, 373–383.
- Segerstrom, S. C., & Miller, G. E. (2004). Psychological stress and the human immune system: A meta-analytic study of 30 years of inquiry. *Psychological Bulletin*, 130, 601–630.
- 1760 Seidlitz, L., Wyer, R. S., & Diener, E. (1997). Cognitive correlates of subjective well-being: The processing of valenced life events by happy and unhappy persons. *Journal of Research in Personality*, 31, 240–256.
- Seligman, M. E., Steen, T. A., Park, N., & Peterson, C. (2005). Positive psychology progress: Empirical validation of interventions. *American Psychologist*, 60, 410–421.
- 1765 Selye, H. (1956). *The stress of life*. New York, NY: McGraw-Hill.
- *Shapiro, D., Jamner, L. D., Goldstein, I. B., & Delfino, R. J. (2001). Striking a chord: Moods, blood pressure, and heart rate in everyday life. *Psychophysiology*, 38, 197–204.
- Sheldon, K. M., & Lyubomirsky, S. (2006). Achieving sustainable gains in happiness: Change your actions, not your circumstances. *Journal of Happiness Studies*, 7, 55–86.
- 1770 Siegler, I. C., Bosworth, H. B., & Elias, M. F. (2003). Adult development and aging. In A. M. Nezu, C. M. Nezu & P. A. Geller (Eds), *Handbook of psychology: Health psychology* (Vol. 9, pp. 487–510). Hoboken, NJ: John Wiley & Sons Inc.
- Siegler, I. C., Poon, L. W., Madden, D. J., & Dilworth-Anderson, P. (2004). Psychological aspects of normal aging. In D. G. Blazer, D. C. Steffens & E. W. Busse (Eds), *The American psychiatric publishing textbook of geriatric psychology* (3rd ed., pp. 121–138). Washington, DC: American Psychiatric Publishing.
- 1775 *Sinha, R., Lovallo, W. R., & Parsons, O. A. (1992). Cardiovascular differentiation of emotions. *Psychosomatic Medicine*, 54, 422–435.
- 1780 Smith, T. W. (2006). Personality as risk and resilience in physical health. *Current Directions in Psychological Science*, 15, 227–231.
- Smith, T. W., & Williams, P. G. (1992). Personality and health: Advantages and limitations of the five-factor model. Special issue: The five-factor model: Issues and applications. *Journal of Personality*, 60, 395–423.
- 1785 Smith, T. W., & Spiro, A. (2002). Personality, health, and aging: Prolegomenon for the next generation. *Journal of Research in Personality*, 36, 363–394.

- Smith, T. W., Glazer, K., Ruiz, J. M., & Gallo, L. C. (2004). Hostility, anger, aggressiveness and coronary heart disease: An interpersonal perspective on personality, emotion, and health. *Journal of Personality, 72*, 1217–1270.
- 1790 *Smyth, J., Ockenfels, M. C., Porter, L., Kirschbaum, C., Hellhammer, D. H., & Stone, A. A. (1998). Stressors and mood measured on a momentary basis are associated with salivary cortisol secretion. *Psychoneuroendocrinology, 23*, 353–370.
- Solomon, G. F., & Benton, D. (1994). Psychoneuroimmunologic aspects of aging. In R. Glaser & J. Kiecolt-Glaser (Eds), *Handbook of human stress and immunity* (pp. 341–363). San Diego, CA: Academic Press.
- 1795 Solomon, G. F., Fiatarone, M. A., Benton, D., Morley, J. E., Bloom, E. T., & Makinodan, T. (1988). Psychoimmunologic and endorphin function in the aged. *Annals of the New York Academy of Science, 521*, 43–58.
- *Steptoe, A., & Holmes, R. (1985). Mood and pulmonary function in adult asthmatics: A pilot self-monitoring study. *British Journal of Medical Psychology, 58*, 87–94.
- 1800 *Steptoe, A., & Wardle, J. (2005). Positive affect and biological function in everyday life. *Neurobiology of Aging. Special Issue: Aging, Diabetes, Obesity. Mood and Cognition, 26*, 108–112.
- *Sternbach, R. A. (1962). Assessing differential autonomic patterns in emotions. *Journal of Psychosomatic Research, 6*, 87–91.
- 1805 *Stevens, M. J., Heise, R. A., & Pfost, K. S. (1989). Consumption of attention versus affect elicited by cognitions in modifying acute pain. *Psychological Reports, 64*, 284–286.
- *Stone, A. A., Cox, D. S., Valdimarsdottir, H., Jandorf, L., & Neale, J. M. (1987). Evidence that secretory IgA antibody is associated with daily mood. *Journal of Personality and Social Psychology, 52*, 988–993.
- 1810 *Stone, A. A., Neale, J. M., Cox, D. S., Napoli, A., Valdimarsdottir, V., & Kennedy-Moore, E. (1994). Daily events are associated with a secretory immune response to an oral antigen in men. *Health Psychology, 13*, 440–446.
- Stones, M. J., & Kozma, A. (1985). Structural relationships among happiness scales: A second order factorial study. *Social Indicators Research, 17*, 19–28.
- 1815 *Stones, M. J., Dornan, B., & Kozma, A. (1989). The prediction of mortality in elderly institution residents. *Journals of Gerontology, 44*, 72–79.
- Strawbridge, W. J., Cohen, R. D., Shema, S. J., & Kaplan, G. A. (1996). Successful aging: Predictors and associated activities. *American Journal of Epidemiology, 144*, 135–141.
- 1820 Suhail, K., & Chaudhry, H. R. (2004). Predictors of subjective well-being in an eastern Muslim culture. *Journal of Social and Clinical Psychology, 23*, 359–376.
- Suls, J., & Bunde, J. (2005). Anger, anxiety, and depression as risk factors for cardiovascular disease: The problems and implications of overlapping affective dimensions. *Psychological Bulletin, 131*, 260–300.
- 1825 *Szczepanski, R., Napolitano, M., Feaganes, J. R., Barefoot, J. C., Luecken, L., Swoap, R. et al. (1997). Relation of mood ratings and neurohormonal responses during daily life in employed women. *International Journal of Behavioral Medicine, 4*, 1–16.
- Thomas, P. D., Goodwin, J. M., & Goodwin, J. S. (1985). Effect of social support on stress-related changes in cholesterol level, uric acid level, and immune function in an elderly sample. *Journal of Psychiatry, 142*, 735–737.
- 1830 *Uchiyama, I. (1992). Differentiation of fear, anger, and joy. *Perceptual and Motor Skills, 74*, 663–667.
- *Uchiyama, I., Hanari, T., Ito, T., Takahashi, K., Okuda, T., Goto, T. et al. (1990). Patterns of psychological and physiological responses for common affects elicited by films. *Psychologia: An International Journal of Psychology in the Orient, 33*, 36–41.
- 1835 *van Domburg, R. T., Schmidt, P. S., van den Brand, M. J. B. M., & Erdman, R. A. M. (2001). Feelings of being disabled as a predictor of mortality in men 10 years after percutaneous coronary transluminal angioplasty. *Journal of Psychosomatic Research, 51*, 469–477.
- *van Eck, M., Berkhof, H., Nicolson, N., & Sulon, J. (1996). The effects of perceived stress, traits, mood states, and stressful daily events on salivary cortisol. *Psychosomatic Medicine, 58*, 447–458.
- 1840 Velten, E. Jr. (1968). A laboratory task for induction of mood states. *Behavior Research and Therapy, 6*, 473–482.

Well-being and health: A meta-analysis 53

- *von Känel, R., Kudielka, B. M., Preckel, D., Hanebuth, D., Herrmann-Lingen, C., Frey, K. et al. (2005). Opposite effect of negative and positive affect on stress procoagulant reactivity. *Physiology & Behavior*, 86, 61–68.
- *von Leupoldt, A., & Dahme, B. (2004). Emotions in a body plethysmograph: Impact of affective film clips on airway resistance. *Journal of Psychophysiology*, 18, 170–176.
- *von Leupoldt, A., & Dahme, B. (2005). Emotions and airway resistance in asthma: Study with whole body plethysmography. *Psychophysiology*, 42, 92–97.
- *Waldstein, S. R., Kop, W. J., Schmidt, L. A., Hauffer, A. J., Krantz, D. S., & Fox, N. A. (2000). Frontal electrocortical and cardiovascular reactivity during happiness and anger. *Biological Psychology*, 55, 3–23.
- Ware, J. E. Jr. (1987). Standards for validating health measures: Definition and content. *Journal of Chronic Diseases*, 40, 473–480.
- Watson, D., & Pennebaker, J. W. (1989). Health complaints, stress, and distress: Exploring the central role of negative affectivity. *Psychological Review*, 96, 234–254.
- Watson, D., & Clark, L. A. (1994). *Manual for the Positive and Negative Affect Schedule (Expanded Form)*. University of Iowa; available from the authors.
- Watson, D., Clark, L. A., & Tellegen, A. (1988). Development and validation of brief measures of positive and negative affect: The PANAS scales. *Journal of Personality and Social Psychology*, 54, 1063–1070.
- *Weaver, J., & Zillmann, D. (1994). Effect on humor and tragedy on discomfort tolerance. *Perceptual & Motor Skills*, 78, 632–634.
- *Weisenberg, M., Raz, T., & Hener, T. (1998). The influence of film-induced mood on pain perception. *Pain*, 76, 365–375.
- Weksler, M. E., & Hausman, P. B. (1982). Effects of aging in the immune response. In D. P. Stites, J. B. Stobo & H. H. Fedenberg (Eds), *Basic and clinical immunology* (pp. 306–313). Los Altos, CA: Lange Medical Publications.
- Westmaas, J. L., Gil-Rivas, V., & Cohen Silver, R. (2007). Designing and implementing interventions to promote health and prevent illness. In H. S. Friedman & R. C. Silver (Eds), *Foundations of health psychology* (pp. 52–70). New York, NY: Oxford University Press.
- *Whorwell, P. J., Houghton, L. A., Taylor, E. E., & Maxton, D. G. (1992). Physiological effects of emotion: Assessment via hypnosis. *Lancet*, 340, 69–72.
- *Wied, M., & Verbatan, M. N. (2001). Affective pictures processing, attention and pain tolerance. *Pain*, 90, 163–172.
- *Williams, J. M., Hogan, T. D., & Andersen, M. B. (1993). Positive states of mind and athletic injury risk. *Psychosomatic Medicine*, 55, 468–472.
- *Wingard, D. L., Berkman, L. F., & Brand, R. J. (1994). A multivariate analysis of health-related practices: A nine-year mortality follow up of the Alameda County Study. In A. Steptoe & J. Wardle (Eds), *Psychosocial processes and health: A reader* (pp. 273–289). New York, NY: Cambridge University Press.
- *Witvliet, C. V., & Vrana, S. R. (1995). Psychophysiological responses as indices of affective dimensions. *Psychophysiology*, 32, 436–443.
- *Worthington, E. L. Jr., & Shumate, M. (1981). Imagery and verbal counseling methods in stress inoculation training for pain control. *Journal of Counseling Psychology*, 28, 1–6.
- *Yogo, Y., Hama, H., Yogo, M., & Matsuyama, Y. (1995). A study of physiological response during emotional imaging. *Perceptual and Motor Skills*, 81, 43–49.
- *Yoshino, S., Fujimori, J., & Kohda, M. (1996). Effects of mirthful laughter on neuroendocrine and immune systems in patients with rheumatoid arthritis. *Journal of Rheumatology*, 23, 793–794.
- *Zachariae, R., Bjerring, P., Zachariae, C., Arendt-Nielsen, L., Nielsen, T., Eldrup, E. et al. (1991). Monocyte chemotactic activity in sera after hypnotically induced emotional states. *Scandinavian Journal of Immunology*, 34, 71–79.
- *Zachariae, R., Jorgensen, M. M., Egekvist, H., & Bjerring, P. (2001). Skin reactions to histamine of healthy subjects after hypnotically induced emotions of sadness, anger, and happiness. *Allergy*, 56, 734–740.
- *Zelman, D. C., Howland, E. W., Nichols, S. N., & Cleeland, C. S. (1991). The effects of induced mood on laboratory pain. *Pain*, 46, 105–111.
- *Zillmann, D., Rockwell, S., Schweitzer, K., & Sundar, S. S. (1993). Does humor facilitate coping with physical discomfort? *Motivation & Emotion*, 17, 1–21.

54 R. T. Howell et al.

- *Zillmann, D., de Wied, M., King-Jablonski, C., & Jenzowsky, S. (1996). Drama-induced affect and pain sensitivity. *Psychosomatic Medicine*, 58, 333–341.
- *Zuckerman, D. M., Kasl, S. V., & Ostfeld, A. M. (1984). Psychosocial predictors of mortality among the elderly poor. The role of religion, well-being, and social contacts. *American Journal of Epidemiology*, 119, 410–423.
- *Zweyer, K., Velker, B., & Ruch, W. (2004). Do cheerfulness, exhilaration, and humor production moderate pain tolerance? A FACS study. *Humor: International Journal of Humor Research*, 17, 85–119.

1905
UNCORRECTED PROOF